



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/authorsrights>

Contents lists available at [SciVerse ScienceDirect](http://www.sciencedirect.com)

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com

Mathematical Analysis/Differential Geometry

Spectral properties of Schrödinger operators on compact manifolds: Rigidity, flows, interpolation and spectral estimates

*Propriétés spectrales d'opérateurs de Schrödinger sur des variétés compactes : Rigidité, flots, interpolation et estimations spectrales*Jean Dolbeault^a, Maria J. Esteban^a, Ari Laptev^b, Michael Loss^c^a Ceremade (UMR CNRS 7534), Université Paris-Dauphine, place du Maréchal-de-Lattre-de-Tassigny, 75775 Paris cedex 16, France^b Department of Mathematics, Imperial College London, Huxley Building, 180 Queen's Gate, SW7 2AZ, UK^c School of Mathematics, Skiles Building, Georgia Institute of Technology, Atlanta, GA 30332-0160, USA

ARTICLE INFO

Article history:

Received 12 March 2013

Accepted 1 July 2013

Available online 31 July 2013

Presented by the Editorial Board

ABSTRACT

This note is devoted to optimal spectral estimates for Schrödinger operators on compact connected Riemannian manifolds without boundary. These estimates are based on the use of appropriate interpolation inequalities and on some recent rigidity results for nonlinear elliptic equations on those manifolds.

© 2013 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Cette note est consacrée à des estimations spectrales optimales pour des opérateurs de Schrödinger sur des variétés riemanniennes compactes et simplement connexes, sans bord. Ces estimations sont basées sur certaines inégalités d'interpolation et sur un résultat récent de rigidité pour des équations elliptiques non linéaires sur ces variétés.

© 2013 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Spectral properties of Schrödinger operators on the sphere

We start by briefly reviewing some results that have been obtained in [4]. Let us define $2^* := \frac{2d}{d-2}$ if $d \geq 3$, and $2^* := \infty$ if $d = 1$ or 2 . We denote by Δ_g the Laplace–Beltrami operator on the unit sphere $\mathbb{S}^d \subset \mathbb{R}^{d+1}$. It is well known (see [2]) that the equation:

$$-\Delta_g u + \frac{\lambda}{q-2} u = u^{q-1}$$

has only constant solutions as long as $q \in (2, 2^*)$ and $\lambda \leq d$. See [3] for a review and various related results. Assume that the measure on \mathbb{S}^d is the one induced by Lebesgue's measure on \mathbb{R}^{d+1} . This convention differs from the one of [4]. The inequality:

$$\|\nabla u\|_{L^2(\mathbb{S}^d)}^2 + \alpha \|u\|_{L^2(\mathbb{S}^d)}^2 \geq \mu(\alpha) \|u\|_{L^q(\mathbb{S}^d)}^2 \quad \forall u \in H^1(\mathbb{S}^d),$$

E-mail addresses: dolbeaul@ceremade.dauphine.fr (J. Dolbeault), esteban@ceremade.dauphine.fr (M.J. Esteban), a.laptev@imperial.ac.uk (A. Laptev), loss@math.gatech.edu (M. Loss).

1631-073X/\$ – see front matter © 2013 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

<http://dx.doi.org/10.1016/j.crma.2013.06.014>

for any $q \in (2, 2^*)$ can be established by standard variational methods. According to [4], the optimal function $\mu : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is concave, increasing, and such that $\mu(\alpha) = \kappa\alpha$ for any $\alpha \leq \frac{d}{q-2}$, $\mu(\alpha) < \kappa\alpha$ for $\alpha > \frac{d}{q-2}$ where $\kappa := |\mathbb{S}^d|^{1-2/q}$ is a normalization factor and:

$$\mu(\alpha) \sim K_{q,d} \alpha^{1-\vartheta} \quad \text{as } \alpha \rightarrow +\infty, \quad \text{where } \vartheta := d \frac{q-2}{2q},$$

$$K_{q,d} := \inf_{v \in H^1(\mathbb{R}^d) \setminus \{0\}} \frac{\|\nabla v\|_{L^2(\mathbb{R}^d)}^2 + \|v\|_{L^2(\mathbb{R}^d)}^2}{\|v\|_{L^q(\mathbb{R}^d)}^2}. \tag{1}$$

Let us define $p = \frac{q}{q-2}$ so that $p \in (1, +\infty)$ if $d = 1$ and $p \in (\frac{d}{2}, +\infty)$ if $d \geq 2$. If we denote by $\mu \mapsto \alpha(\mu)$ the inverse function of $\alpha \mapsto \mu(\alpha)$ and by $\lambda_1(-\Delta_g - V)$ the lowest (nonpositive) eigenvalue of $-\Delta_g - V$, then we have the estimate

$$|\lambda_1(-\Delta_g - V)| \leq \alpha(\|V\|_{L^p(\mathbb{S}^d)}) \quad \forall V \in L^p(\mathbb{S}^d),$$

for any nonnegative $V \in L^p(\mathbb{S}^d)$. Moreover we have $\alpha(\mu)^{p-d/2} = L_{p-\frac{d}{2},d}^1 \mu^p (1 + o(1))$ as $\mu \rightarrow +\infty$ where $L_{\gamma,d}^1 := (K_{q,d})^{-(\gamma+d/2)}$. Equality is achieved for any $\mu > 0$ by some nonnegative V , which is constant if and only if $\mu \leq \frac{d}{2}(p-1)$.

The case $q \in (1, 2)$ can also be covered and we refer to [4] for further details. This case leads to estimates from below for the first eigenvalue of the operator $-\Delta_g + W$, where W is a positive potential.

2. A rigidity result on compact manifolds and a subcritical interpolation inequality

From here on we shall assume that (\mathfrak{M}, g) is a smooth compact connected Riemannian manifold of dimension $d \geq 1$, without boundary, and let Δ_g be the Laplace–Beltrami operator on \mathfrak{M} . We shall denote by dv_g the volume element and by \mathfrak{R} the Ricci tensor. Let λ_1 be the lowest positive eigenvalue of $-\Delta_g$.

For such manifolds, a new rigidity result has been recently established in [5], thus extending a series of results of [6,2,9,1,8]. In order to state this result, let us define the quantities:

$$\theta = \frac{(d-1)^2(p-1)}{d(d+2)+p-1} \quad \text{and} \quad Q_g u := H_g u - \frac{g}{d} \Delta_g u - \frac{(d-1)(p-1)}{\theta(d+3-p)} \left[\frac{\nabla u \otimes \nabla u}{u} - \frac{g}{d} \frac{|\nabla u|^2}{u} \right]$$

where $H_g u$ denotes Hessian of u , and

$$\Lambda_\star := \inf_{u \in H^2(\mathfrak{M}) \setminus \{0\}} \frac{(1-\theta) \int_{\mathfrak{M}} (\Delta_g u)^2 dv_g + \frac{\theta d}{d-1} \int_{\mathfrak{M}} [\|Q_g u\|^2 + \mathfrak{R}(\nabla u, \nabla u)] dv_g}{\int_{\mathfrak{M}} |\nabla u|^2 dv_g}. \tag{2}$$

It is not difficult to see that $\Lambda_\star \leq \lambda_1$.

Theorem 1. (Cf. [5, Theorem 3].) Assume that Λ_\star is strictly positive. Then for any $q \in (1, 2) \cup (2, 2^*)$ and any $\lambda \in (0, \Lambda_\star)$, the equation:

$$-\Delta_g v + \frac{\lambda}{q-2} (v - v^{q-1}) = 0$$

has 1 as its unique positive solution in $C^2(\mathfrak{M})$.

Note that in the particular case $\mathfrak{M} = \mathbb{S}^d$, $\Lambda_\star = \lambda_1(-\Delta_g) = d$. The proof relies on the nonlinear flow

$$u_t = u^{2-2\beta} \left(\Delta_g u + (1 + \beta(q-2)) \frac{|\nabla u|^2}{u} \right), \quad \beta = \frac{(d+2)(d+3-p)\theta}{(d-1)^2(p-1)^2 - (d+2)^2(p-2)\theta}, \tag{3}$$

that can also be used to prove the following A - B -type interpolation inequality (see [2,7]). Let us define:

$$\kappa := \text{vol}_g(\mathfrak{M})^{1-2/q}.$$

Theorem 2. (Cf. [5, Theorem 4].) For any $q \in (1, 2) \cup (2, 2^*)$ if $d \geq 3$, $q \in (1, 2) \cup (2, \infty)$ if $d = 1$ or 2 , the inequality:

$$\|\nabla v\|_{L^2(\mathfrak{M})}^2 \geq \frac{\Lambda}{q-2} [\kappa \|v\|_{L^q(\mathfrak{M})}^2 - \|v\|_{L^2(\mathfrak{M})}^2] \quad \forall v \in H^1(\mathfrak{M})$$

holds for some optimal $\Lambda \in [\Lambda_\star, \lambda_1]$ if $\Lambda_\star > 0$. Moreover, if $\Lambda_\star < \lambda_1$, then we have $\Lambda_\star < \Lambda \leq \lambda_1$.

The above results hold true because the flow (3) contracts:

$$\mathcal{F}[u] := \int_{\mathfrak{M}} |\nabla(u^\beta)|^2 dv_g + \frac{\Lambda_\star}{q-2} \left[\int_{\mathfrak{M}} u^{2\beta} dv_g - \kappa \left(\int_{\mathfrak{M}} u^{\beta q} dv_g \right)^{2/q} \right].$$

The above choices for θ and β are optimal for this contraction property: see [5].

As a consequence and exactly as in the case of the sphere, we get the first result of this note.

Proposition 3. Assume that $q \in (2, 2^\star)$ if $d \geq 3$, or $q \in (2, \infty)$ if $d = 1$ or 2 . There exists a concave increasing function $\mu : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that $\mu(\alpha) = \kappa \alpha$ for any $\alpha \leq \frac{\Lambda_\star}{q-2}$, $\mu(\alpha) < \kappa \alpha$ for $\alpha > \frac{\Lambda_\star}{q-2}$ and

$$\|\nabla u\|_{L^2(\mathfrak{M})}^2 + \alpha \|u\|_{L^q(\mathfrak{M})}^2 \geq \mu(\alpha) \|u\|_{L^q(\mathfrak{M})}^2 \quad \forall u \in H^1(\mathfrak{M}).$$

The asymptotic behavior of μ is given by $\mu(\alpha) \sim K_{q,d} \alpha^{1-\vartheta}$ as $\alpha \rightarrow +\infty$, with $\vartheta = d \frac{q-2}{2q}$ and $K_{q,d}$ defined by (1).

Proof. There is an optimal function for the interpolation inequality, as can be shown by standard variational techniques. Applying Theorem 1 to the solutions of the Euler–Lagrange equations completes the proof for fixed values of α . As an infimum on u of affine functions with respect to α , the function $\alpha \rightarrow \mu(\alpha)$ is concave. It remains to establish the properties of α for large values of μ .

Using a well-chosen test function obtained by scaling an optimal function for (1) on the tangent plane to an arbitrary point of \mathfrak{M} , one can prove that $\limsup_{\alpha \rightarrow +\infty} \alpha^{\vartheta-1} \mu(\alpha) \leq K_{q,d}$. Arguing by contradiction as in [4, Proposition 10], we can find a sequence $(\alpha_n)_{n \in \mathbb{N}}$ such that $\lim_{n \rightarrow +\infty} \alpha_n = +\infty$ and $\lim_{n \rightarrow +\infty} \alpha_n^{\vartheta-1} \mu(\alpha_n) < K_{q,d}$, and a sequence of optimal functions $(u_n)_{n \in \mathbb{N}}$ such that $\|u_n\|_{L^q(\mathbb{S}^d)} = 1$, which concentrates because $\limsup_{n \rightarrow +\infty} \alpha_n^{\vartheta} \|u_n\|_{L^2(\mathbb{S}^d)}^2 < K_{q,d}$. Some classical surgery and a convexity inequality provide a contradiction by constructing a minimizing sequence for (1). \square

3. Ground state estimates for Schrödinger operators on Riemannian manifolds

In this section, we keep using the notations of Section 2 and generalize to (\mathfrak{M}, g) the spectral results established for the sphere in [4]. By inverting the function $\alpha \mapsto \mu(\alpha)$, we see that $\alpha : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is increasing, convex and satisfies: $\alpha(\mu) = \frac{\mu}{\kappa}$ for any $\mu \in (0, \frac{\kappa \Lambda_\star}{q-2})$, $\alpha(\mu) > \frac{\mu}{\kappa}$ for $\mu \in (\frac{\kappa \Lambda_\star}{q-2}, +\infty)$. With $L_{\gamma,d}^1 := (K_{q,d})^{-p}$, $\gamma = p - \frac{d}{2}$, we obtain for a general manifold \mathfrak{M} the same behavior of $\mu \mapsto \alpha(\mu)$ when $\mu \rightarrow +\infty$ as in the case of a sphere.

Let $\mu := \|V\|_{L^p(\mathbb{S}^d)}$. Since p and $\frac{q}{2}$ are Hölder conjugate exponents, it follows from Hölder's inequality that:

$$\int_{\mathfrak{M}} |\nabla u|^2 dv_g - \int_{\mathfrak{M}} V|u|^2 dv_g + \alpha(\mu) \int_{\mathfrak{M}} |u|^2 dv_g \geq \|\nabla u\|_{L^2(\mathbb{S}^d)}^2 - \mu \|u\|_{L^q(\mathbb{S}^d)}^2 + \alpha(\mu) \|u\|_{L^2(\mathbb{S}^d)}^2$$

with equality if V^{p-1} and $|u|^2$ are proportional. The right-hand side is nonnegative according to Proposition 3. By taking the infimum of the left-hand side, we can deduce an estimate of the lowest, nonpositive eigenvalue $\lambda_1(-\Delta_g - V)$ of $-\Delta_g - V$, which provides us with our first main result.

Theorem 4. Let $d \geq 1$, $p \in (1, +\infty)$ if $d = 1$ and $p \in (\frac{d}{2}, +\infty)$ if $d \geq 2$ and assume that $\Lambda_\star > 0$. With the above notations and definitions, for any nonnegative $V \in L^p(\mathfrak{M})$, we have:

$$|\lambda_1(-\Delta_g - V)| \leq \alpha(\|V\|_{L^p(\mathfrak{M})}). \tag{4}$$

Moreover, we have $\alpha(\mu)^{p-\frac{d}{2}} = L_{p-\frac{d}{2},d}^1 \mu^p (1 + o(1))$ as $\mu \rightarrow +\infty$.

The estimate (4) is optimal in the sense that for any $\mu \in (0, +\infty)$, there exists a nonnegative function V such that $\mu = \|V\|_{L^p(\mathfrak{M})}$ and $|\lambda_1(-\Delta_g - V)| = \alpha(\mu)$. Moreover, if $\mu < \frac{\kappa \Lambda_\star}{q-2}$, $\alpha(\mu) = \frac{\mu}{\kappa}$ and equality in (4) is achieved by constant potentials.

In the case of operators $-\Delta_g + W$ on \mathfrak{M} , where W is a nonnegative potential, following again the same arguments as in [4] in the case of the sphere, and the rigidity result of Theorem 1, we obtain our second main result.

Theorem 5. Let $d \geq 1$, $p \in (0, +\infty)$. There exists an increasing concave function $v : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, satisfying $v(\beta) = \beta/\kappa$, for any $\beta \in (0, \frac{p+1}{2} \kappa \Lambda)$ if $p > 1$, such that for any positive potential W we have:

$$\lambda_1(-\Delta + W) \geq v(\beta) \quad \text{with} \quad \beta = \left(\int_{\mathfrak{M}} W^{-p} dv_g \right)^{1/p}.$$

Moreover, for large values of β , we have $v(\beta)^{-(p+\frac{d}{2})} = L_{-(p+\frac{d}{2}),d}^1 \beta^{-p} (1 + o(1))$ as $\beta \rightarrow +\infty$.

With $p = \frac{q}{2-q}$, the spectral estimate of [Theorem 5](#) is derived from the interpolation inequality:

$$\|\nabla u\|_{L^2(\mathfrak{M})}^2 + \beta \left(\int_{\mathfrak{M}} |u|^q dv_g \right)^{2/q} \geq \nu(\beta) \|u\|_{L^2(\mathfrak{M})}^2 \quad \forall u \in H^1(\mathfrak{M}).$$

The concentration phenomena leading to the asymptotics for large norms of W can be studied as in [Proposition 3](#): see [\[4\]](#) for the proof in the case of a sphere. We omit the details of the proof of [Theorem 5](#).

Acknowledgements

J.D. and M.J.E. have been partially supported by ANR grants *CBDif* and *NoNAP*. They thank the Mittag-Leffler Institute, where part of this research was carried out, for hospitality. M.L. was supported in part by NSF grant DMS-0901304.

References

- [1] D. Bakry, M. Ledoux, Sobolev inequalities and Myers's diameter theorem for an abstract Markov generator, *Duke Math. J.* 85 (1996) 253–270.
- [2] M.-F. Bidaut-Véron, L. Véron, Nonlinear elliptic equations on compact Riemannian manifolds and asymptotics of Emden equations, *Invent. Math.* 106 (1991) 489–539.
- [3] J. Dolbeault, M.J. Esteban, M. Kowalczyk, M. Loss, Sharp interpolation inequalities on the sphere: New methods and consequences, *Chin. Ann. Math. Ser. B* 34 (2013) 99–112.
- [4] J. Dolbeault, M.J. Esteban, A. Laptev, Spectral estimates on the sphere, *Anal. Partial Differ. Equ.* (2013), in press, preprint hal-00770755.
- [5] J. Dolbeault, M.J. Esteban, M. Loss, Nonlinear flows and rigidity results on compact manifolds, preprint hal-00784887.
- [6] B. Gidas, J. Spruck, Global and local behavior of positive solutions of nonlinear elliptic equations, *Commun. Pure Appl. Math.* 34 (1981) 525–598.
- [7] E. Hebey, *Nonlinear Analysis on Manifolds: Sobolev Spaces and Inequalities*, Courant Lecture Notes in Mathematics, vol. 5, New York University Courant Institute of Mathematical Sciences, New York, 1999.
- [8] J.R. Licois, L. Véron, A class of nonlinear conservative elliptic equations in cylinders, *Ann. Sc. Norm. Sup. Pisa Cl. Sci. (4)* 26 (1998) 249–283.
- [9] J.R. Licois, L. Véron, Un théorème d'annulation pour des équations elliptiques non linéaires sur des variétés riemanniennes compactes, *C. R. Acad. Sci. Paris, Ser. I* 320 (1995) 1337–1342.