DOI: 10.1002/mana.202000196

ORIGINAL PAPER

MATHEMATISCHE NACHRICHTEN

Sharp bounds for eigenvalues of biharmonic operators with complex potentials in low dimensions

Orif O. Ibrogimov¹ | David Krejčiřík¹ | Ari Laptev^{2,3}

¹ Department of Mathematics, Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Trojanova 13, Prague 2 12000, Czechia

² Department of Mathematics, Imperial College London, Huxley Building, 180 Queen's Gate, London SW7 2AZ, UK

³ Sirius Mathematical Center, Sirius University of Science and Technology, 1 Olympic Ave, Sochi 354340, Russia

Correspondence

David Krejčiřík, Department of Mathematics, Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Trojanova, 13, Prague 212000, Czechia. Email: david.krejcirik@fjfi.cvut.cz

· - ·

Funding information Grantová Agentura České Republiky, Grant/Award Number: 20-17749X

Abstract

We derive sharp quantitative bounds for eigenvalues of biharmonic operators perturbed by complex-valued potentials in dimensions one, two and three.

KEYWORDS

biharmonic operators, bounds for eigenvalues, complex potentials, Birman–Schwinger principle

MSC (2020) 35J30, 35P15, 81Q12

1 | INTRODUCTION

Spectral theory of self-adjoint operators has exhibited an enormous development since the discovery of quantum mechanics in the beginning of the last century and by now it can be regarded as well understood in many respects. Recent years have brought new motivations for considering *non-self-adjoint* operators, too, including quantum mechanics again, but this theory is still in its infancy.

A strong impetus for a systematic study of spectral properties of Schrödinger operators with complex-valued potentials goes back to the celebrated result of Davies et al. in 2001 [1] showing that every discrete eigenvalue of the one-dimensional operator $-\Delta \dot{+}V$ lies in the closed disk of the complex plane centred at the origin and with the radius equal to $\frac{1}{4} ||V||_{L^1(\mathbb{R})}^2$. This bound is sharp in the sense that there are potentials with eigenvalues lying on the boundary circle. The purpose of this paper is to extend this type of bounds to biharmonic operators

$$H_V := \Delta^2 + V \quad \text{in} \quad L^2(\mathbb{R}^d) \tag{1.1}$$

in dimensions d = 1, 2, 3. The feature of our results is that our bounds are sharp and quantitative (at least if d = 1, 3). Moreover, potentials of minimal regularity are considered and we cover embedded eigenvalues, too.

The pioneering work [1] has been followed by extensive investigations devoted to bounds on the number and magnitudes of eigenvalues of Schrödinger operators with complex-valued potentials. Instead of listing the huge number of articles by

various groups on operators not dealt with in this paper, we refer to the recent work [13] containing a rather extensive bibliography. On the other hand, related issues for higher-order partial differential operators seem to be just scarcely considered in the literature, even in the self-adjoint setting. The reader is referred to [7, 9, 10, 17, 20, 22, 26]; see also [4–6, 11, 14] for other spectral questions.

We recall the importance of biharmonic operators in continuum mechanics and in particular linear elasticity theory and the solution of Stokes flows. To the best of our knowledge, the only work where eigenvalue bounds for non-selfadjoint biharmonic operators were considered previously is the recent paper of Enblom [10]. We also refer to recent Hulko's paper [17] for bounds on the number of discrete eigenvalues of biharmonic operators perturbed by exponentially decaying complex-valued potentials. While the author of [10] proceeds in a rather extensive generality (polyharmonic operators in any dimension), her bounds are not quantitative (in the sense that the involved constants depend on the arguments of the individual eigenvalues under consideration) and embedded eigenvalues are not covered. For these reasons, it is our belief that our results are of interest even if restricted to biharmonic operators in low dimensions only. Moreover, our methodology in principle enables one to obtain analogous results also for higher-order polyharmonic operators.

Before stating our main results, we remark that the operator (1.1) is introduced in a standard way as an m-accretive operator obtained as a form sum of the bi-Laplacian $H_0 := \Delta^2$ with domain $H^4(\mathbb{R}^d)$ and a relatively form-bounded potential $V : \mathbb{R}^d \to \mathbb{C}$. We refer to Section 2 for more details. More specifically, our standing assumption is that there exist numbers $a \in (0, 1)$ and $b \in \mathbb{R}$ such that, for all $\psi \in H^2(\mathbb{R}^d)$,

$$\int_{\mathbb{R}^d} |V| |\psi|^2 \le a \int_{\mathbb{R}^d} |\Delta \psi|^2 + b \int_{\mathbb{R}^d} |\psi|^2.$$
(1.2)

In particular, potentials $V \in L^1(\mathbb{R}^d)$ are covered by this hypothesis. Furthermore, in the same way, it is also possible to proceed in a greater generality and give a meaning to the distributional Dirac delta potential δ , which is explicitly solvable. Putting $\|\delta\|_{L^1(\mathbb{R}^d)} := 1$ by convention, Theorem 1.1 remains valid in this more general setting.

Our first result is a biharmonic analogue of the celebrated result of [1] for one-dimensional Schrödinger operators.

Theorem 1.1. Let $d \in \{1, 2, 3\}$ and assume that $V \in L^1(\mathbb{R}^d)$. Then there exists a universal constant $C_d > 0$ such that

$$\sigma_{\mathbf{p}}(H_V) \subset \left\{ \lambda \in \mathbb{C} : |\lambda| \le C_d \|V\|_{L^1(\mathbb{R}^d)}^{4/(4-d)} \right\}.$$
(1.3)

Moreover, one can take $C_1 = \frac{1}{4}$ *and* $C_3 = \frac{1}{4} \frac{1}{(4\pi)^4}$ *for dimensions* d = 1 *and* d = 3*, respectively.*

The theorem is sharp due to the weak-coupling asymptotics

$$\inf \sigma \left(H_{\beta V} \right) = -c_d \, \beta^{4/(4-d)} \left\| V \right\|_{L^1(\mathbb{R}^d)}^{4/(4-d)} + o \left(\beta^{4/(4-d)} \right) \quad \text{as} \quad \beta \to 0^+$$

valid for every sufficiently regular, non-positive, real-valued potential *V* with $c_1 := \frac{1}{4}$, $c_2 := \frac{1}{64}$ and $c_3 := \frac{1}{4} \frac{1}{(4\pi)^4}$, cf. [22, Eq. (8)]. We do not know whether one can take $C_2 = \frac{1}{64}$ in (1.3) for dimension d = 2, but it is natural to conjecture so. The result is sharp also in the sense that for d = 1 (respectively, d = 3) one has $\sigma_p(H_{\alpha\delta}) = \left\{\frac{1}{4}\alpha^{4/3}e^{-i\pi/3}\right\}$ for every $\alpha \in \mathbb{C}$ such that $\Re \alpha < |\Im \alpha|$ (respectively, $\sigma_p(H_{4\pi\alpha\delta}) \supset \left\{-\frac{1}{4}\alpha^4\right\}$ if $\Re \alpha < -|\Im \alpha|$).

In the next series of results, we go beyond the L^1 -potentials in the three-dimensional case. First, we employ the Rollnik class of potentials (cf. [24, Ch. I]), which consist of all $V \in L^1_{loc}(\mathbb{R}^3)$ such that

$$\|V\|_{R(\mathbb{R}^3)}^2 := \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{|V(x)| |V(y)|}{|x - y|^2} \, dx \, dy < \infty.$$
(1.4)

In Section 2, we argue that $V \in R(\mathbb{R}^3)$ is a sufficient condition to guarantee (1.2), so H_V is well defined.

IBROGIMOV ET AL.

Theorem 1.2. Let d = 3 and assume that $V \in R(\mathbb{R}^3)$. Then

$$\sigma_{\mathrm{p}}(H_V) \subset \left\{ \lambda \in \mathbb{C} : |\lambda| \le \frac{1}{2} \frac{\|V\|_{R(\mathbb{R}^3)}^2}{(4\pi)^2} \right\}.$$
(1.5)

This result has an important consequence for potentials belonging to $L^{3/2}(\mathbb{R}^3) \hookrightarrow R(\mathbb{R}^3)$. In fact, using a sharp version of the Hardy–Littlewood–Sobolev inequality (cf. [21, Thm. 4.3]) which quantifies the embedding, the bound (1.5) immediately implies the following result.

Corollary 1.3. Let d = 3 and assume that $V \in L^{3/2}(\mathbb{R}^3)$. Then

$$\sigma_{\rm p}(H_V) \subset \left\{ \lambda \in \mathbb{C} : \ |\lambda| \le \frac{1}{8} \frac{1}{(4\pi)^{2/3}} \|V\|_{L^{3/2}(\mathbb{R}^3)}^2 \right\}.$$
(1.6)

In view of Theorem 1.1 and Corollary 1.3, complex interpolation applied to an appropriate analytic family of Birman–Schwinger type operators yields the following explicit eigenvalue bound.

Theorem 1.4. Let d = 3 and assume that $V \in L^p(\mathbb{R}^3)$ for some $p \in (1, 3/2)$. Then

$$\sigma_{p}(H_{V}) \subset \left\{ \lambda \in \mathbb{C} : |\lambda| \leq 2^{\frac{12-14p}{4p-3}} (4\pi)^{\frac{4p-8}{4p-3}} \|V\|_{L^{p}(\mathbb{R}^{3})}^{\frac{4p}{4p-3}} \right\}.$$
(1.7)

Finally, we establish the following robust result.

Theorem 1.5. Let d = 3 and assume (1.2). Then

$$\sigma_{\mathrm{p}}(H_{V}) \subset \left\{ \lambda \in \mathbb{C} : |\lambda| \leq \left(\sup_{\substack{\psi \in H^{1}(\mathbb{R}^{3}) \\ \psi \neq 0}} \frac{\int_{\mathbb{R}^{3}} |V| |\psi|^{2}}{\int_{\mathbb{R}^{3}} |\nabla \psi|^{2}} \right)^{2} \right\}.$$
(1.8)

2

This theorem is particularly useful to cover the Hardy-type potentials $x \mapsto |x|^{-2}$, which belong neither to $L^1(\mathbb{R}^3)$ nor $R(\mathbb{R}^3)$. For this reason, let us consider the class of potentials, which consist of all $V \in L^1_{loc}(\mathbb{R}^3)$ such that

$$\|V\|_{H(\mathbb{R}^3)} := \operatorname{ess\,sup}_{x \in \mathbb{R}^3} |x|^2 |V(x)| < \infty.$$
(1.9)

Employing the classical Hardy inequality

$$\int_{\mathbb{R}^3} |\nabla \psi|^2 \ge \frac{1}{4} \int_{\mathbb{R}^3} \frac{|\psi(x)|^2}{|x|^2} \, dx \tag{1.10}$$

valid for every $\psi \in H^1(\mathbb{R}^3)$, it is easy to check that $V \in H(\mathbb{R}^3)$ is a sufficient condition to guarantee (1.2). Furthermore, using (1.10) in (1.8), we get the following immediate consequence.

Corollary 1.6. Let d = 3 and assume that $V \in H(\mathbb{R}^3)$. Then

$$\sigma_{\mathbf{p}}(H_V) \subset \left\{ \lambda \in \mathbb{C} : |\lambda| \le 16 \|V\|_{H(\mathbb{R}^3)}^2 \right\}.$$
(1.11)

In this paper we leave aside dimension d = 4, which exhibits the same type of difficulties as dimension two for Schrödinger operators. The reason for explicit constants in our one- and three-dimensional theorems is due to the

availability of explicit forms of the resolvent kernels of H_0 in these dimensions. On the other hand, in view of the subcriticality of H_0 in dimensions $d \ge 5$, a different kind of results is expected in analogy with the case of Schrödinger operators (cf. [13, 15, 19]): the point spectrum of H_V should be empty for all potentials which are sufficiently small in a suitable sense. This type of results are also left open in this paper.

The organisation of the paper is as follows. In Section 2 we rigorously introduce the biharmonic operators (1.1) and determine sufficient conditions on the potential *V* which guarantee the basic hypothesis (1.2). In Section 3 we establish the main tool of our approach, namely a sort of the Birman–Schwinger principle covering embedded eigenvalues, too. The proofs of Theorems 1.1–1.5 are given in Section 4. Finally, in Section 5 we introduce explicitly solvable models in terms of Dirac delta potentials and argue about the optimality of Theorem 1.1 for dimensions d = 1 and d = 3.

2 | DEFINITION OF BIHARMONIC OPERATORS

First of all, let us comment on the definition of H_V given in (1.1).

Let H_0 be the self-adjoint operator in $L^2(\mathbb{R}^d)$ associated with the quadratic form

$$h_0[\psi] := \int_{\mathbb{R}^d} |\Delta \psi|^2, \quad \mathsf{D}(h_0) := H^2(\mathbb{R}^d).$$
(2.1)

One has $D(H_0) = H^4(\mathbb{R}^d)$ and $H_0 = \Delta^2$. The spectrum of H_0 is purely absolutely continuous and coincides with the semiaxis $[0, +\infty)$. If $d \le 4$, the operator H_0 is *critical* in the sense that $\inf \sigma(H_0 + V) < 0$ whenever $V \in C_0^{\infty}(\mathbb{R}^d)$ is real-valued, non-positive and non-trivial. On the other hand, if $d \ge 5$, the operator is *subcritical* in the sense that its spectrum is stable under the non-positive perturbations above; more specifically, H_0 satisfies a Hardy-type inequality $H_0 \ge \rho$ in the sense of forms, where $\rho : \mathbb{R}^d \to [0, \infty)$ is non-trivial.

Let v be a quadratic form in $L^2(\mathbb{R}^d)$, which is relatively bounded with respect to h_0 with the relative bound less than one. That is, $D(v) \supset H^2(\mathbb{R}^d)$ and there exist numbers $a \in (0, 1)$ and $b \in \mathbb{R}$ such that, for all $\psi \in H^2(\mathbb{R}^d)$,

$$|v[\psi]| \le a \int_{\mathbb{R}^d} |\Delta \psi|^2 + b \int_{\mathbb{R}^d} |\psi|^2.$$
(2.2)

Then the sum $h_V := h_0 + v$ is a closed sectorial form with $D(H_V) = H^2(\mathbb{R}^d)$, which gives rise to an m-sectorial operator H_V in $L^2(\mathbb{R}^d)$ via the representation theorem (cf. [18, Thm. VI.2.1]).

For example, if $V \in L^1_{loc}(\mathbb{R}^d)$ is such that

$$v[\psi] := \int_{\mathbb{R}^d} V |\psi|^2, \quad \mathsf{D}(v) := \bigg\{ \psi \in L^2\big(\mathbb{R}^d\big) \, : \, \int_{\mathbb{R}^d} |V| |\psi|^2 < \infty \bigg\},$$

verifies (2.2) (which coincides with (1.2) in this case), then we write $H_V = H_0 + V$ as in (1.1) and understand the plus symbol in the sense of forms described above. It is important to keep in mind that H_V may differ from the operator sum $H_0 + V$. Since the adjoint operator satisfies $H_V^* = H_V = \mathcal{T} H_V \mathcal{T}$, where \mathcal{T} is the complex conjugation operator defined by $\mathcal{T}\psi := \overline{\psi}, H_V$ is a \mathcal{T} -self-adjoint operator. Consequently, the residual spectrum of H_V is always empty (cf. [8, Sec. III.5]). Furthermore, if V vanishes at infinity in a suitable sense, then (any kind of) the essential spectrum of H_V coincides with the semi-axis $[0, +\infty)$.

Let us now discuss sufficient conditions which guarantee (2.2).

By the Sobolev embedding theorem ([3, Thm. 5.4]), every function $\psi \in H^2(\mathbb{R}^d)$ is bounded and continuous as long as $d \leq 3$. More specifically (cf. [23, Theorem IX.28]), for any positive α there is $\beta \in \mathbb{R}$ such that, for all $\psi \in H^2(\mathbb{R}^d)$,

$$\|\psi\|_{L^{\infty}(\mathbb{R}^d)} \le \alpha \|\Delta\psi\|_{L^2(\mathbb{R}^d)} + \beta \|\psi\|_{L^2(\mathbb{R}^d)}.$$
(2.3)

Consequently, any potential $V \in L^1(\mathbb{R}^d) + L^{\infty}(\mathbb{R}^d)$ satisfies (1.2) with the relative bound equal to zero (i.e. *a* can be chosen arbitrarily small). Furthermore, the inequality (2.3) enables us to give a meaning to the operator $H_{\delta} = H_0 + \delta$, where δ is the Dirac delta function, by setting $v[\psi] := |\psi(0)|^2$, $\mathsf{D}(v) := H^2(\mathbb{R}^d)$.

Rollnik potentials are relatively form-bounded with respect to the Laplacian with the relative bound equal to zero (cf. [23, Theorem X.19]). More specifically, if $V \in R(\mathbb{R}^3)$, then for any positive α there is $\beta \in \mathbb{R}$ such that, for all $\psi \in H^1(\mathbb{R}^3)$,

$$\int_{\mathbb{R}^3} |V| |\psi|^2 \leq \alpha \|\nabla \psi\|_{L^2(\mathbb{R}^3)}^2 + \beta \|\psi\|_{L^2(\mathbb{R}^3)}^2.$$

Consequently, if additionally $\psi \in H^2(\mathbb{R}^3)$, then

$$\int_{\mathbb{R}^{3}} |V| |\psi|^{2} \leq \alpha (\psi, -\Delta \psi)_{L^{2}(\mathbb{R}^{3})} + \beta \|\psi\|_{L^{2}(\mathbb{R}^{3})}$$

$$\leq \alpha \|\psi\|_{L^{2}(\mathbb{R}^{3})} \|\Delta \psi\|_{L^{2}(\mathbb{R}^{3})} + \beta \|\psi\|_{L^{2}(\mathbb{R}^{3})}^{2}.$$
(2.4)

From this inequality, it is easy to conclude that any potential $V \in R(\mathbb{R}^3) + L^{\infty}(\mathbb{R}^3)$ satisfies (1.2) with the relative bound equal to zero.

Finally, if $V \in H(\mathbb{R}^3)$, then the Hardy inequality (1.10) yields, for all $\psi \in H^1(\mathbb{R}^3)$,

$$\int_{\mathbb{R}^3} |V| |\psi|^2 \le \|V\|_{H(\mathbb{R}^3)} \int_{\mathbb{R}^3} \frac{|\psi(x)|^2}{|x|^2} \, dx \le 4 \|V\|_{H(\mathbb{R}^3)} \|\nabla\psi\|_{L^2(\mathbb{R}^3)}^2.$$

Hence *V* is form-subordinated with respect to the Laplacian. Proceeding as in (2.4), we conclude that any potential $V \in H(\mathbb{R}^3) + L^{\infty}(\mathbb{R}^3)$ satisfies (1.2) with the relative bound equal to zero.

3 | THE BIRMAN-SCHWINGER PRINCIPLE

The main role in our proof of Theorems 1.1-1.5 is played by the Birman-Schwinger operator

$$K_{z} := |V|^{1/2} (H_{0} - z)^{-1} V_{1/2} \quad \text{with} \quad V_{1/2} := |V|^{1/2} \operatorname{sgn}(V),$$
(3.1)

where sgn : $\mathbb{C} \to \mathbb{C}$ is the complex signum function defined by

$$\operatorname{sgn}(z) := \begin{cases} \frac{z}{|z|} & \text{if } z \neq 0, \\ 0 & \text{if } z = 0. \end{cases}$$

We abuse the notation by using the same symbols for maximal operators of multiplication and their generating functions. The operator K_z is well defined (on its natural domain of the composition of three operators) for all $z \in \mathbb{C}$ and $d \ge 1$.

If $z \notin [0, +\infty)$, however, K_z is a bounded operator under our hypothesis (1.2). Indeed, $V_{1/2}$ maps $L^2(\mathbb{R}^d)$ to $H^{-2}(\mathbb{R}^d)$ by duality, $(H_0 - z)^{-1}$ is an isomorphism between $H^{-2}(\mathbb{R}^d)$ and $H^2(\mathbb{R}^d)$, and the latter space is mapped by $|V|^{1/2}$ back to $L^2(\mathbb{R}^d)$. Furthermore, we have a useful formula for the integral kernel of K_z :

$$K_z(x,y) = |V|^{1/2}(x) \widetilde{G}_z(x,y) V_{1/2}(y), \qquad (3.2)$$

where \tilde{G}_z is the *Green's function* of $H_0 - z$, i.e. the integral kernel of the resolvent $(H_0 - z)^{-1}$. What is more, for dimensions d = 1 and d = 3, we have explicit formulae for \tilde{G}_z , whereas the latter can be expressed in terms of modified Hankel functions for d = 2. In fact, using the identity

$$\left(\Delta^2 - z\right)^{-1} = \frac{1}{2k} \left[(-\Delta - k)^{-1} - (-\Delta + k)^{-1} \right],\tag{3.3}$$

1338 | MATHEMATISCHE NACHRICHTEN

where $k \in \mathbb{C} \setminus [0, \infty)$ is such that $k^2 = z$ (throughout the paper we choose the principal branch of the square root), and the well-known formulae for the integral kernels of the resolvent of the Laplacian in these dimensions yield

$$\widetilde{G}_{z}(x,y) = \frac{1}{2k} [G_{k}(x,y) - G_{-k}(x,y)],$$
(3.4)

where

$$G_{k}(x,y) := \begin{cases} \frac{e^{-\sqrt{-k}|x-y|}}{2\sqrt{-k}} & \text{if } d = 1, \\ \frac{1}{2\pi} K_{0} \left(\sqrt{-k}|x-y|\right) & \text{if } d = 2, \\ \frac{1}{4\pi} \frac{e^{-\sqrt{-k}|x-y|}}{|x-y|} & \text{if } d = 3. \end{cases}$$
(3.5)

Here K_0 is a modified Bessel (also called Macdonald's) function (see [2, Sec. 9] or [16, Sec. 8.4–8.5]).

The following lemma provides an integral criterion for the existence of solutions to the differential eigenvalue equation corresponding to H_V and can be considered as a one-sided version of the conventional Birman–Schwinger principle extended to possible eigenvalues embedded in $[0, +\infty)$ as well. Its proof is inspired by the ones of the similar results in [12, 13]. By $\varphi \in L_0^2(\mathbb{R}^d)$ in the lemma below we mean $\varphi \in L^2(\mathbb{R}^d)$ and that supp φ is compact.

Lemma 3.1. Let $d \in \{1, 2, 3\}$ and assume (1.2). If $H_V \psi = \lambda \psi$ with some $\lambda \in \mathbb{C}$ and $\psi \in D(H_V)$, then $\phi := |V|^{1/2} \psi \in L^2(\mathbb{R}^d)$ obeys

$$\forall \varphi \in L_0^2(\mathbb{R}^d), \quad \lim_{\varepsilon \to 0^{\pm}} \left(\varphi, K_{\lambda + i\varepsilon} \phi\right) = -(\varphi, \phi).$$
(3.6)

Proof. First, we notice that (1.2) implies that $\phi \in L^2(\mathbb{R}^d)$ whenever $\psi \in D(H_V) \subset H^2(\mathbb{R}^d)$. Let $\varphi \in L^2_0(\mathbb{R}^d)$ be fixed. Given any $\lambda \in \mathbb{C}$, there is $\varepsilon_0 > 0$ such that $\lambda + i\varepsilon \notin [0, +\infty)$ for all real ε satisfying $0 < |\varepsilon| < \varepsilon_0$. We have

$$(\varphi, K_{\lambda+i\varepsilon}\phi) = \iint_{\mathbb{R}^d \times \mathbb{R}^d} \overline{\varphi(x)} |V|^{1/2}(x) G_{\lambda+i\varepsilon}(x, y) V(y) \psi(y) dx dy$$

$$= \int_{\mathbb{R}^d} \eta_{\varepsilon}(y) V(y) \psi(y) dy,$$

$$(3.7)$$

where

$$\eta_{\varepsilon} := \int_{\mathbb{R}^d} \overline{\varphi(x)} \, |V|^{1/2}(x) \, G_{\lambda+i\varepsilon}(x,\cdot) \, dx = \left(H_0 - \lambda - i\varepsilon\right)^{-1} |V|^{1/2} \, \overline{\varphi} \,,$$

where the second equality holds due to the symmetry $G_z(x, y) = G_z(y, x)$. In view of (1.2), $|V|^{1/2}\overline{\varphi} \in L^2(\mathbb{R}^d)$. Since $\varepsilon \neq 0$ is so small that $\lambda + i\varepsilon \notin \sigma(H_0)$, we have $\eta_{\varepsilon} \in D(H_0) = H^4(\mathbb{R}^d)$. In particular, $\eta_{\varepsilon} \in H^2(\mathbb{R}^d)$ and the weak formulation of the eigenvalue equation $H_V \psi = \lambda \psi$ yields

$$\begin{split} \int_{\mathbb{R}^d} \eta_{\varepsilon}(y) V(y) \psi(y) \, dy &= -\left(\Delta \overline{\eta_{\varepsilon}}, \Delta \psi\right) + \lambda \left(\overline{\eta_{\varepsilon}}, \psi\right) \\ &= -\left(\Delta \overline{\psi}, \Delta \eta_{\varepsilon}\right) + \lambda \left(\overline{\psi}, \eta_{\varepsilon}\right) \\ &= -\left(\Delta \overline{\psi}, \Delta \eta_{\varepsilon}\right) + \left(\lambda + i\varepsilon\right) \left(\overline{\psi}, \eta_{\varepsilon}\right) - i\varepsilon \left(\overline{\psi}, \eta_{\varepsilon}\right) \\ &= -\left(\overline{\psi}, |V|^{1/2} \overline{\psi}\right) - i\varepsilon \left(\overline{\psi}, \eta_{\varepsilon}\right) \\ &= -\left(\varphi, |V|^{1/2} \psi\right) - i\varepsilon \left(\overline{\eta_{\varepsilon}}, \psi\right) \,. \end{split}$$
(3.8)

Here the last but one equality follows from the weak formulation of the resolvent equation $(H_0 - \lambda - i\varepsilon)\eta_{\varepsilon} = |V|^{1/2}\overline{\varphi}$. Consequently, (3.7) and (3.8) imply (3.6) after taking the limit $\varepsilon \to 0^{\pm}$, provided that $\varepsilon (\bar{\eta}_{\varepsilon}, \psi) \to 0$ as $\varepsilon \to 0$. To see the latter, we write

$$\left|\left(\overline{\eta_{\varepsilon}},\psi
ight)
ight|=\left|\left(arphi,M_{arepsilon}\psi
ight)
ight|\leq\left\|arphi
ight\|\left\|M_{arepsilon}
ight\|\left\|\psi
ight\|,$$

where $M_{\varepsilon} := \chi_{\Omega} |V|^{1/2} (H_0 - \lambda - i\varepsilon)^{-1}$ with $\Omega := \operatorname{supp} \varphi$, and it remains to show that $\varepsilon ||M_{\varepsilon}||$ tends to zero as $\varepsilon \to 0$. Following [24, Thm. III.6], we use the resolvent kernels (3.4) with (3.5) and the estimate $||M_{\varepsilon}|| \le ||M_{\varepsilon}||_{HS}$.

For d = 1, we have

$$\begin{split} \|M_{\varepsilon}\|_{\mathrm{HS}}^{2} &= \frac{1}{4|k|^{2}} \iint_{\Omega \times \mathbb{R}} |V(x)| \, |G_{k}(x,y) - G_{-k}(x,y)|^{2} dx \, dy \\ &\leq \frac{1}{2|k|^{2}} \iint_{\Omega \times \mathbb{R}} |V(x)| \left[\frac{e^{-2\Re \sqrt{-k} |x-y|}}{4|k|} + \frac{e^{-2\Re \sqrt{k} |x-y|}}{4|k|} \right] dx \, dy \\ &= \frac{1}{8|k|^{3}} \left[\frac{1}{\Re \sqrt{-k}} + \frac{1}{\Re \sqrt{k}} \right] \int_{\Omega} |V(x)| dx \,, \end{split}$$

where the last integral is bounded because $V \in L^1_{\text{loc}}(\mathbb{R})$ as a consequence of (1.2). Elementary calculations show that $|k|^3 \Re \sqrt{\pm k}$ can not decay faster than $|\varepsilon|^{7/4}$ as $\varepsilon \to 0$. Hence, we have $||M_{\varepsilon}|| = \mathcal{O}(|\varepsilon|^{-7/8})$ as $\varepsilon \to 0$, which concludes the proof of the lemma for d = 1.

For d = 3, we have analogous estimates

$$\begin{split} \|M_{\varepsilon}\|_{\mathrm{HS}}^{2} &\leq \frac{1}{32\pi^{2}|k|^{2}} \iint_{\Omega \times \mathbb{R}^{3}} |V(x)| \left[\frac{e^{-2\Re\sqrt{-k}|x-y|}}{|x-y|^{2}} + \frac{e^{-2\Re\sqrt{k}|x-y|}}{|x-y|^{2}} \right] dx \, dy \\ &= \frac{1}{16\pi |k|^{2}} \left[\frac{1}{\Re\sqrt{-k}} + \frac{1}{\Re\sqrt{k}} \right] \int_{\Omega} |V(x)| dx \, . \end{split}$$

Since $|k|^2 \Re \sqrt{\pm k}$ can not decay faster than $|\varepsilon|^{5/4}$ as $\varepsilon \to 0$, we have $||M_{\varepsilon}|| = \mathcal{O}(|\varepsilon|^{-5/8})$ as $\varepsilon \to 0$. This concludes the proof of the lemma for d = 3.

For d = 2, we need to do rather involved estimates. Using the integral representation of the Macdonald's function (cf. [16, § 8.432]), we can write Green's function for the biharmonic operator as

$$\widetilde{G}_{z}(x,y) = \frac{1}{4\pi k} \int_{1}^{\infty} \frac{e^{-\sqrt{-k}|x-y|t} - e^{-\sqrt{k}|x-y|t}}{\sqrt{t^{2} - 1}} dt.$$

Next, we observe that, for t > 1,

$$\int_0^\infty \left| e^{-\sqrt{-kts}} + e^{-\sqrt{kts}} \right|^2 ds \le 2 \int_0^\infty \left(e^{-2\Re\sqrt{-kts}} + e^{-2\Re\sqrt{kts}} \right) ds = \frac{1}{t} \left[\frac{1}{\Re\sqrt{k}} + \frac{1}{\Re\sqrt{-k}} \right].$$

In view of this estimate, the Minkowski inequality yields

$$\begin{split} \|M_{\varepsilon}\|_{\mathrm{HS}}^{2} &\leq \frac{1}{8\pi^{2}|k|^{2}} \bigg(\int_{\Omega} |V(x)| \, dx \bigg) \bigg(\int_{1}^{\infty} \bigg(\int_{0}^{\infty} \left| e^{-\sqrt{-k}ts} + e^{-\sqrt{k}ts} \right|^{2} ds \bigg)^{1/2} \frac{dt}{\sqrt{t^{2}-1}} \bigg)^{2} \\ &= \frac{1}{8\pi^{2}|k|^{2}} \left[\frac{1}{\Re\sqrt{-k}} + \frac{1}{\Re\sqrt{k}} \right] \bigg(\int_{\Omega} |V(x)| \, dx \bigg) \bigg(\int_{1}^{\infty} \frac{1}{\sqrt{t}\sqrt{t^{2}-1}} \, dt \bigg)^{2}. \end{split}$$

We already know that $|k|^2 \Re \sqrt{\pm k}$ can not decay faster than $|\varepsilon|^{5/4}$ as $\varepsilon \to 0$. Hence, again we have $||M_{\varepsilon}|| = \mathcal{O}(|\varepsilon|^{-5/8})$ as $\varepsilon \to 0$, which concludes the proof of the lemma for d = 2.

The preceding lemma can be viewed as a precise statement of one side of the Birman–Schwinger principle under the minimal regularity assumption (1.2) on the potential. It roughly says that if λ is an eigenvalue of H_V , then -1 is an eigenvalue of an integral equation related to K_{λ} . If $\lambda \notin \sigma(H_0)$ the converse implication also holds, but it is not generally true if $\lambda \in \sigma(H_0)$ (cf. [24, Sec. III.2]) and it is not needed for the purpose of this paper. In fact, we exclusively use the following corollary of Lemma 3.1.

Corollary 3.2. Let $d \in \{1, 2, 3\}$ and assume (1.2). Let $\lambda \in \mathbb{C}$ be arbitrary. If either $\lambda \in \rho(H_0)$ and $||K_{\lambda}|| < 1$ or $\lambda \in \sigma(H_0)$ and $\lim \inf_{\varepsilon \to 0^{\pm}} ||K_{\lambda+i\varepsilon}|| < 1$, then $\lambda \notin \sigma_p(H_V)$.

Proof. Let $\lambda \in \sigma_p(H_V)$, let ψ be a corresponding eigenfunction and set $\phi := |V|^{1/2} \psi$. If it were the case that $\phi = 0$, then the definition of H_V would yield

$$(\varphi, (H_0 - \lambda)\psi) = (\varphi, (H_V - \lambda)\psi) - \int_{\mathbb{R}^d} \overline{\varphi} V\psi = 0,$$

for all $\varphi \in H^2(\mathbb{R}^d)$, and consequently, $H_0 \psi = \lambda \psi$. Unless $\psi = 0$, this would mean that λ is an eigenvalue of H_0 , which is impossible. Hence, we conclude that $\phi \neq 0$.

Next, if $\lambda \in \rho(H_0)$, then Lemma 3.1 with $\varepsilon = 0$ implies

$$\|\phi\|^2 \|K_{\lambda}\| \ge |(\phi, K_{\lambda}\phi)| = \|\phi\|^2$$
(3.9)

and thus $||K_{\lambda}|| \ge 1$.

If $\lambda \in \sigma(H_0)$, we set $\phi_n := \xi_n \phi$ for every positive *n*, where $\xi_n(x) := \xi(x/n)$ and $\xi \in C_0^{\infty}(\mathbb{R}^d)$ is a usual cut-off function satisfying $\xi(x) = 1$ for $|x| \le 1$ and $\xi(x) = 0$ for $|x| \ge 2$. As in (3.9), we have

$$\|\phi_n\|\|\phi\|\|K_{\lambda+i\varepsilon}\| \ge |(\phi_n, K_{\lambda+i\varepsilon}\phi)|$$

In view of $\phi_n \in L^2_0(\mathbb{R}^d)$, we can invoke Lemma 3.1 and take the limit $\varepsilon \to 0^{\pm}$ to conclude

$$\|\phi_n\| \|\phi\| \liminf_{\varepsilon \to 0^{\pm}} \|K_{\lambda+i\varepsilon}\| \ge |(\phi_n, \phi)|$$

The desired claim now follows by taking the limit $n \to \infty$.

4 | PROOFS

First, we establish two elementary inequalities which will be crucial in obtaining sharp eigenvalue bounds.

Lemma 4.1. For all nonnegative real numbers p and q, the following inequality holds:

$$\frac{1}{2}(e^{q-p} + e^{p-q}) \le e^{p+q} - \sin(p+q).$$
(4.1)

Proof. Let us rewrite (4.1) as

$$e^{-2p} + e^{-2q} + 2e^{-(p+q)}\sin(p+q) \le 2.$$
(4.2)

1340 MAIH

 \square

Since (4.2) is symmetric with respect to *p* and *q*, there is no loss of generality in assuming that $0 \le p \le q$. Let us fix $p = p_0 \ge 0$ and analyse the smooth function

$$\Phi(q) := e^{-2p_0} + e^{-2q} + 2e^{-(p_0+q)}\sin(p_0+q)$$

on the interval $[p_0, \infty)$. It is easy to check that every possible critical point q_0 of Φ must satisfy the equation

$$e^{q_0-p_0} = \cos(p_0+q_0) - \sin(p_0+q_0).$$

In view of this observation, some elementary calculations yield that

$$\Phi(q_0) = 2e^{-2p_0} \left(1 - \sin^2(p_0 + q_0) \right) \le 2.$$
(4.3)

On the other hand, we have

$$\Phi(p_0) = 2e^{-2p_0} (1 + \sin(2p_0)) \le 2, \quad \lim_{q \uparrow +\infty} \Phi(q) = e^{-2p_0} \le 1, \tag{4.4}$$

the first estimate being the consequence of the elementary inequality $sin(t) \le t \le e^t - 1$ for $t \ge 0$. Hence, we conclude from (4.3) and (4.4) that

$$\max_{q\geq p_0}\Phi(q)\leq 2,$$

which proves the claim in (4.2).

Lemma 4.2. For all nonnegative real numbers p and q, the following inequality holds:

$$\frac{1}{2}(e^{q-p} + e^{p-q}) \le (p^2 + q^2)e^{p+q} + \cos(p+q).$$
(4.5)

Proof. Let us rewrite (4.5) as

$$e^{-2p} + e^{-2q} - 2e^{-(p+q)}\cos(p+q) \le 2(p^2 + q^2).$$
(4.6)

Since (4.6) is symmetric with respect to p and q, there is no loss of generality in assuming that $0 \le p \le q$. Let us fix $p = p_0 \ge 0$ and analyse the smooth function

$$\Phi(q) := e^{-2p_0} + e^{-2q} - 2e^{-(p_0+q)}\cos(p_0+q) - 2(p_0^2+q^2)$$

on the interval $[p_0, \infty)$. We have

$$\Phi'(q) = -2e^{-2q} + 2e^{-(p_0+q)}\cos(p_0+q) + 2e^{-(p_0+q)}\sin(p_0+q) - 4q.$$
(4.7)

On the other hand, for all $s, t \in \mathbb{R}$, we have the well-known inequalities $e^t \ge \cos(t) + \sin(t)$ and $e^s \ge 1 + s$. Applying these inequalities with $t = p_0 + q$ and s = -2q, we obtain

$$\Phi'(q) \le -2e^{-2q} + 2 - 4q \le 0. \tag{4.8}$$

Therefore, Φ is non-increasing on $[0, \infty)$ and thus we have

$$\Phi(q) \le \Phi(p_0) = 2e^{-2p_0} + 2e^{-2p_0}\cos(2p_0) - 4p_0^2 \le 0,$$

for all $q \ge p_0$, where for the last step one needs to recall the inequality

$$1 - \cos(t) \le \frac{1}{2}t^2 \le \frac{1}{2}t^2e^t, \quad t \ge 0.$$

Hence, for each fixed $p \ge 0$, we have $\Phi(q) \le 0$ for all $q \ge p$. This proves the desired claim.

Remark 4.3. We note that the expressions on the left-hand sides of (4.2) and (4.6) are nonnegative for all $p, q \ge 0$. In the same way as in Lemma 4.1 one can show that

$$\frac{1}{2}(e^{q-p} + e^{p-q}) \le e^{p+q} + \cos(p+q)$$
(4.9)

holds for all $p, q \ge 0$.

Now we are in a position to establish our theorems.

4.1 | Proof of Theorem 1.1

We start with the case $\lambda \in \mathbb{C} \setminus [0, \infty)$ and denote by *k* the principal square root of λ as before. Throughout the proof we assume the parameter *k* to be fixed.

First, for each $d \in \{1, 2, 3\}$, we justify the existence of a universal constant $c_d > 0$ such that the corresponding Green's function of the biharmonic operators obeys the following pointwise estimate

$$\left|\widetilde{G}_{\lambda}(x,y)\right| \le \frac{c_d}{|k|^{2-d/2}}.$$
(4.10)

For the case d = 1, elementary calculations show that the inequality

$$\left|\sqrt{k}\,e^{-\sqrt{-k}\,|x-y|} - \sqrt{-k}\,e^{-\sqrt{k}\,|x-y|}\right| \le \sqrt{2}\sqrt{|k|}$$

is equivalent to (4.1) with $p = \Re \sqrt{k} |x - y| \ge 0$ and $q = \Im \sqrt{k} |x - y| \ge 0$ if $\arg(\lambda) \in (0, \pi]$ or with $p = \Re \sqrt{-k} |x - y| \ge 0$ and $q = \Im \sqrt{-k} |x - y| \ge 0$ if $\arg(\lambda) \in (-\pi, 0)$. Hence, we have (4.10) with the constant $c_1 := \frac{1}{2\sqrt{2}}$, i.e.

$$\left|\widetilde{G}_{\lambda}(x,y)\right| \leq \frac{1}{2\sqrt{2}|k|^{3/2}}.$$

For the case d = 3, elementary calculations show that the inequality

$$\left| e^{-\sqrt{-k} |x-y|} - e^{-\sqrt{k} |x-y|} \right| \le \sqrt{2}\sqrt{|k|} |x-y|$$

is equivalent to (4.5) with $p = \Re \sqrt{k} |x - y| \ge 0$ and $q = \Im \sqrt{k} |x - y| \ge 0$ if $\arg(\lambda) \in (0, \pi]$ or with $p = \Re \sqrt{-k} |x - y| \ge 0$ and $q = \Im \sqrt{-k} |x - y| \ge 0$ if $\arg(\lambda) \in (-\pi, 0)$. That is why we have (4.10) with the constant $c_3 := \frac{1}{4\sqrt{2\pi}}$, i.e.

$$\left|\widetilde{G}_{\lambda}(x,y)\right| \leq \frac{1}{4\sqrt{2}\pi\sqrt{|k|}}$$

For the case d = 2, on the account of the asymptotic expansion of the Macdonald's function $K_0(\zeta)$ for small ζ , we can write (cf. [16, § 8.447])

$$G_k(x,y) = -\frac{1}{2\pi} \ln \sqrt{-k} - \frac{1}{2\pi} \left(\gamma + \ln \frac{|x-y|}{2} \right) + o(|x-y|), \quad |x-y| \to 0^+,$$

where γ is the Euler constant. Hence, it follows that

$$G_k(x,y) - G_{-k}(x,y) = \frac{1}{2\pi} \left(\ln \sqrt{k} - \ln \sqrt{-k} \right) + o(|x-y|)$$
$$= -\frac{i}{4} + o(|x-y|), \quad |x-y| \to 0^+,$$

and consequently, recalling (3.4), we get

$$\widetilde{G}_{\lambda}(x,y) = \frac{i}{8\pi k} \operatorname{Arg}(k) + o(|x-y|), \quad |x-y| \to 0^+.$$

On the other hand, in view of the well-known asymptotic expansion of the Macdonald's function $K_0(\zeta)$ for large ζ , we have

$$G_k(x,y) = \sqrt{\frac{\pi}{2\sqrt{-k}|x-y|}} e^{-\sqrt{-k}|x-y|} (1+\mathcal{O}(1)) = \mathcal{O}\left(\frac{1}{\sqrt[4]{|x-y|}}\right), \quad |x-y| \to +\infty,$$

and consequently,

$$\widetilde{G}_{\lambda}(x,y) = \mathcal{O}(1), \quad |x-y| \to +\infty.$$

Since these asymptotic expansions are uniform in k and the Macdonald's function is entire on the right half-plane, we conclude the existence of a universal constant $c_2 > 0$ satisfying (4.10).

In view of (4.10), now we can estimate the norm of the Birman–Schwinger operator as follows

$$\|K_{\lambda}\|^{2} \leq \|K_{\lambda}\|_{\mathrm{HS}}^{2} = \iint_{\mathbb{R}^{d} \times \mathbb{R}^{d}} |V(x)| \left| \widetilde{G}_{\lambda}(x, y) \right|^{2} |V(y)| \, dx \, dy \leq \frac{c_{d}^{2}}{|k|^{4-d}} \|V\|_{L^{1}(\mathbb{R}^{d})}^{2} \tag{4.11}$$

and thus

$$\|K_{\lambda}\| \le \frac{c_d \|V\|_{L^1(\mathbb{R}^d)}}{|\lambda|^{1-d/4}}.$$
(4.12)

If $\lambda \in (0, \infty)$, then the same analysis applied for $\lambda + i\varepsilon$ with $\varepsilon > 0$ (instead of λ) yields

$$\liminf_{\varepsilon \to 0^{\pm}} \|K_{\lambda+i\varepsilon}\| \le \liminf_{\varepsilon \to 0^{\pm}} \frac{c_d \|V\|_{L^1(\mathbb{R}^d)}}{|\lambda+i\varepsilon|^{1-d/4}} = \frac{c_d \|V\|_{L^1(\mathbb{R}^d)}}{|\lambda|^{1-d/4}}.$$

Hence, Corollary 3.2 implies that $\lambda \in \mathbb{C} \setminus \{0\}$ cannot be an eigenvalue for H_V unless it holds that

$$c_d \|V\|_{L^1(\mathbb{R}^d)} \ge |\lambda|^{1-d/4}.$$

This completes the proof since the origin $\lambda = 0$ obviously belongs to the right-hand side of (1.3).

1344 MATHEMATISCHE NACHRICHTEN –

4.2 | Proof of Theorem 1.2

First, assume that $\lambda \in \mathbb{C} \setminus [0, \infty)$ and let *k* be the principal square root of λ . Elementary calculations show that the estimate

$$\left|e^{-\sqrt{-k}|x-y|} - e^{-\sqrt{k}|x-y|}\right| \le \sqrt{2}$$

is equivalent to (4.9) with $p = \Re \sqrt{k} |x - y| \ge 0$, $q = \Im \sqrt{k} |x - y| \ge 0$ if $\arg(\lambda) \in (0, \pi]$, and with $p = \Re \sqrt{-k} |x - y| \ge 0$, $q = \Im \sqrt{-k} |x - y| \ge 0$ if $\arg(\lambda) \in (-\pi, 0)$. Hence, the Green's function can be estimated as

$$\left|\widetilde{G}_{\lambda}(x,y)\right| \leq \frac{1}{4\sqrt{2\pi|k||x-y|}}$$

for all distinct $x, y \in \mathbb{R}^3$. Using this, we can estimate the norm of the Birman–Schwinger operator in terms of the Rollnik norm of *V* as follows

$$\|K_{\lambda}\|^{2} \leq \|K_{\lambda}\|_{\mathrm{HS}}^{2} = \iint_{\mathbb{R}^{3} \times \mathbb{R}^{3}} |V(x)| \left|\widetilde{G}_{\lambda}(x, y)\right|^{2} |V(y)| \, dx \, dy \leq \frac{1}{32\pi^{2} |k|^{2}} \|V\|_{R(\mathbb{R}^{3})}^{2}.$$

Hence, we have

$$\|K_{\lambda}\| \le \frac{1}{4\pi\sqrt{|\lambda|}} \frac{\|V\|_{R(\mathbb{R}^{3})}}{\sqrt{2}}.$$
 (4.13)

If $\lambda \in (0, \infty)$, then the same analysis applied for $\lambda + i\varepsilon$ with $\varepsilon > 0$ (instead of λ) yields

$$\liminf_{\varepsilon \to 0^{\pm}} \left\| K_{\lambda + i\varepsilon} \right\| \le \liminf_{\varepsilon \to 0^{\pm}} \frac{1}{4\pi \sqrt{|\lambda + i\varepsilon|}} \frac{\|V\|_{R(\mathbb{R}^3)}}{\sqrt{2}} = \frac{1}{4\pi \sqrt{|\lambda|}} \frac{\|V\|_{R(\mathbb{R}^3)}}{\sqrt{2}}$$

Hence, Corollary 3.2 implies that $\lambda \in \mathbb{C} \setminus \{0\}$ cannot be an eigenvalue for H_V if

$$\frac{1}{4\pi} \frac{\|V\|_{R(\mathbb{R}^3)}}{\sqrt{2}} < \sqrt{|\lambda|}$$

This completes the proof since $\lambda = 0$ obviously belongs to the right-hand side of (1.5).

4.3 | Proof of Theorem 1.4

Let $\lambda \in \mathbb{C} \setminus [0, \infty)$ be fixed and suppose that 1 . Define the operator family

$$T_{\lambda}(\zeta) := |V|^{(-\zeta/3+1)p/2} (H_0 - \lambda)^{-1} |V|^{(-\zeta/3+1)p/2},$$

for $\zeta \in \mathbb{C}$ with $0 \leq \Re \zeta \leq 1$. By the hypothesis, we have $|V|^p \in L^1(\mathbb{R}^3)$. Hence, we deduce from (4.12) that

$$\left\|T_{\lambda}(iy)\right\| \le \left\||V|^{p/2} \left(H_0 - \lambda\right)^{-1} |V|^{p/2}\right\| \le \frac{1}{4\sqrt{2\pi}} \frac{\|V\|_{L^p(\mathbb{R}^3)}^p}{|\lambda|^{1/4}},\tag{4.14}$$

IBROGIMOV ET AL.

for all $y \in \mathbb{R}$. On the other hand, we also have that $|V|^{2p/3} \in L^{3/2}(\mathbb{R}^3) \subset R(\mathbb{R}^3)$. Consequently, for all $y \in \mathbb{R}$, first using the bound (4.13) and then the Hardy–Littlewood–Sobolev inequality (cf. [21, Thm. 4.3]), we obtain

$$\begin{split} \left\| T_{\lambda}(1+iy) \right\| &\leq \left\| |V|^{p/3} \left(H_0 - \lambda \right)^{-1} |V|^{p/3} \right\| \leq \frac{1}{4\pi\sqrt{2}} \frac{\left\| |V|^{2p/3} \right\|_{R(\mathbb{R}^3)}}{\sqrt{|\lambda|}} \\ &\leq \frac{1}{2\sqrt{2|\lambda|} (4\pi)^{1/3}} \left\| |V|^{2p/3} \right\|_{L^{3/2}(\mathbb{R}^3)} = \frac{1}{2\sqrt{2} (4\pi)^{1/3}} \frac{\left\| V \right\|_{L^p(\mathbb{R}^3)}^{2p/3}}{\sqrt{|\lambda|}}. \end{split}$$
(4.15)

Next, we note that the operator family $T_{\lambda}(\cdot)$ is analytic in the strip $0 < \Re \zeta < 1$, continuous and uniformly bounded in its closure $0 \le \Re \zeta \le 1$, the latter observation being an immediate consequence of the estimates (4.14) and (4.15). Hence, Stein's complex interpolation theorem (cf. [25, Thm. 1]) applies and yields

$$\|T_{\lambda}(\theta)\| \leq \frac{1}{2^{\frac{1}{2}+\theta}} \frac{1}{(4\pi)^{1-\frac{2}{3}\theta}} \frac{\|V\|_{L^{p}(\mathbb{R}^{3})}^{\left(1-\frac{2}{3}\right)p}}{|\lambda|^{\frac{1}{4}+\frac{\theta}{4}}},$$

for all $\theta \in [0, 1]$. Now Corollary 3.2 completes the proof for non-embedded eigenvalues, if we choose $\theta = 3 - 3/p \in (0, 1)$ and observe that $||K_{\lambda}|| = ||T_{\lambda}(3 - 3/p)||$. If $\lambda \in (0, \infty)$, then the same analysis applied for $\lambda + i\varepsilon$ with $\varepsilon > 0$ instead of λ as in the proofs of Theorems 1.1, 1.2 yields the result.

4.4 | Proof of Theorem 1.5

First, assume that $\lambda \in \mathbb{C} \setminus [0, \infty)$ and let *k* be the principal square root of λ . By the triangle inequality, we have

$$\|K_{\lambda}\| \leq \frac{1}{2|k|} \left[\left\| |V|^{1/2} (-\Delta - k)^{-1} V_{1/2} \right\| + \left\| |V|^{1/2} (-\Delta + k)^{-1} V_{1/2} \right\| \right],$$

where the integral kernel reads now (cf. (3.5))

$$(-\Delta - k)^{-1}(x, y) = \frac{e^{-\sqrt{-k}|x-y|}}{4\pi |x-y|}$$

Using the pointwise estimate

$$|(-\Delta - k)^{-1}(x, y)| \le (-\Delta)^{-1}(x, y)$$

valid for all $x, y \in \mathbb{R}$ with $x \neq y$ and $k \in \mathbb{C}$, we estimate as in [13, proof of Lem. 1]

$$\left\| |V|^{1/2} (-\Delta \pm k)^{-1} V_{1/2} \right\| \le \left\| |V|^{1/2} (-\Delta)^{-1} |V|^{1/2} \right\| \le \left\| |V|^{1/2} (-\Delta)^{-1/2} \right\| \left\| (-\Delta)^{-1/2} |V|^{1/2} \right\|.$$

Finally, noticing that

$$\left\| (-\Delta)^{-1/2} |V|^{1/2} \right\| = \left\| |V|^{1/2} (-\Delta)^{-1/2} \right\| = \left(\sup_{\substack{\psi \in H^1(\mathbb{R}^3) \\ \psi \neq 0}} \frac{\int_{\mathbb{R}^3} |V| |\psi|^2}{\int_{\mathbb{R}^3} |\nabla \psi|^2} \right)^{1/2},$$

we arrive at the estimate

1346

$$\left\|K_{\lambda}\right\| \leq \frac{1}{\sqrt{|\lambda|}} \left(\sup_{\substack{\psi \in H^{1}(\mathbb{R}^{3})\\\psi \neq 0}} \frac{\int_{\mathbb{R}^{3}} |V| |\psi|^{2}}{\int_{\mathbb{R}^{3}} |\nabla \psi|^{2}} \right).$$
(4.16)

We notice as before that the origin $\lambda = 0$ trivially satisfies the estimate in (1.8). If $\lambda \in (0, \infty)$, then by repeating the analysis for $\lambda + i\varepsilon$ with $\varepsilon > 0$ (instead of λ) we see that the right-hand side of (4.16) dominates $\lim \inf_{\varepsilon \to 0^{\pm}} ||K_{\lambda+i\varepsilon}||$. The claimed inclusion thus immediately follows from Corollary 3.2.

5 | OPTIMALITY OF THE BOUNDS

In this section we discuss two concrete examples which demonstrate the optimality of the constants corresponding to the eigenvalue bounds for dimensions d = 1 and d = 3 in Theorem 1.1. Both examples are given in terms of distributional Dirac delta potentials, but an approximation by regular potentials is also mentioned. We do not consider the delta potential in two dimensions because the constant C_2 of Theorem 1.1 is not explicit (nonetheless, preliminary computations confirm that the weak-coupling constant $c_2 = \frac{1}{64}$ is achieved by the Dirac delta potential also for d = 2).

5.1 | The delta potential in one dimension

As described in Section 2, the operator $H_{\alpha\delta} = H_0 + \alpha\delta$ in $L^2(\mathbb{R})$ with $\alpha \in \mathbb{C}$ should be understood as the operator associated with the quadratic form

$$h_{\alpha\delta}[\psi] := \int_{\mathbb{R}} |\psi''|^2 + \alpha |\psi(0)|^2, \quad \mathsf{D}(h_{\alpha\delta}) := H^2(\mathbb{R}).$$

One has

$$H_{\alpha\delta}\psi(x) = \psi^{\prime\prime\prime\prime}(x), \quad x \in \mathbb{R} \setminus \{0\},$$

$$\mathsf{D}(H_{\alpha\delta}) = \left\{\psi \in H^4(\mathbb{R} \setminus \{0\}) \cap H^3(\mathbb{R}) : \psi^{\prime\prime\prime}(0^+) - \psi^{\prime\prime\prime}(0^-) = -\alpha\,\psi(0)\right\}.$$
(5.1)

Note that the function values at 0 are well defined due to the Sobolev embedding $H^4(\mathbb{R}) \hookrightarrow C^3(\mathbb{R})$. Since the perturbation is a point interaction, it follows that $\sigma_{ess}(H_{\alpha\delta}) = [0, +\infty)$.

Let us look for eigenvalues of $H_{\alpha\delta}$, i.e. we consider $H_{\alpha\delta}\psi = k^4\psi$ with the convention

$$\Re k > 0 \quad \text{and} \quad \Im k > 0.$$
 (5.2)

The general solution of the differential equation $\psi^{\prime\prime\prime\prime} = k^4 \psi$ in any open real interval reads

$$\psi(x) = C_1 e^{kx} + C_2 e^{-kx} + C_3 e^{ikx} + C_4 e^{-ikx}, \qquad (5.3)$$

where C_1, C_2, C_3, C_4 are arbitrary complex numbers. Considering these general solution in the separate intervals $(-\infty, 0)$ and $(0, +\infty)$, and imposing the integrability requirements $(C_1, C_4 = 0 \text{ if } x > 0 \text{ and } C_2, C_3 = 0 \text{ if } x < 0)$ together with the interface conditions at zero due to (5.1) (continuity up to the second derivative and the jump condition for the third derivative), we arrive at the condition

$$k^3 = \frac{1}{2^{3/2}} \, \alpha \, e^{-i\pi/4}$$

In order to make this equation compatible with (5.2), we see that the point spectrum of $H_{\alpha\delta}$ is not empty if, and only if, arg $\alpha \in (\pi/4, 7\pi/4)$, or equivalently, $\Re \alpha < |\Im \alpha|$. If this condition is satisfied, $H_{\alpha\delta}$ possesses one eigenvalue corresponding to the right-hand side put to the power 4/3. Summing up,

1347

$$\sigma_{\mathbf{p}}(H_{\alpha\delta}) = \begin{cases} \left\{ \frac{1}{4} \,\alpha^{4/3} \, e^{-i\pi/3} \right\} & \text{if} \quad \Re\alpha < |\Im\alpha| \,, \\ \\ \emptyset & \text{if} \quad \Re\alpha \ge |\Im\alpha| \,. \end{cases}$$

Notice that the arguments of α for which the point spectrum is empty correspond to real points intersecting the half-axis $[0, +\infty)$. So any eigenvalue of $H_{\alpha\delta}$, if it exists, is necessarily discrete.

In particular, choosing $\alpha := e^{i\theta}$ and varying $\theta \in (\pi/4, 7\pi/4)$, any boundary point of the closed disk $\{\lambda \in \mathbb{C} : |\lambda| \le \frac{1}{4}\}$ except for the point $\lambda = \frac{1}{4}$ will be an eigenvalue of $H_{\alpha\delta}$. This proves the optimality of Theorem 1.1 for d = 1 because $\|\alpha\delta\|_{L^1(\mathbb{R})} = |\alpha| = 1$.

Remark 5.1. One can also justify the optimality of the constant in the eigenvalue bound corresponding to the inclusion (1.3) for d = 1 in the limit $\varepsilon \to 0^+$ by considering the following family of regular but singularly scaled potentials

$$\delta_{\varepsilon}(x) := \begin{cases} \frac{1}{\varepsilon} & \text{if } |x| < \varepsilon/2, \\ 0 & \text{otherwise.} \end{cases}$$

Of course $\delta_{\varepsilon} \to \delta$ in the sense of distributions as $\varepsilon \to 0^+$. What is more, the point spectrum of $H_{\alpha\delta_{\varepsilon}}$ converges to the point spectrum of $H_{\alpha\delta}$ as $\varepsilon \to 0^+$. This can be checked by explicitly solving the differential equations of the eigenvalue problem in the separate intervals where δ_{ε} is constant and matching the solutions at the boundary points.

5.2 | The delta potential in three dimensions

In analogy with the one-dimensional situation above, we consider the operator $H_{4\pi\alpha\delta} = H_0 + 4\pi\alpha\delta$ in $L^2(\mathbb{R}^3)$ with $\alpha \in \mathbb{C}$ associated with the quadratic form

$$h_{4\pi\alpha\delta}[\psi] := \int_{\mathbb{R}^3} |\Delta\psi|^2 + 4\pi\alpha \, |\psi(0)|^2 \,, \quad \mathsf{D}\big(h_{4\pi\alpha\delta}\big) := H^2\big(\mathbb{R}^3\big) \,.$$

Again, $\sigma_{\text{ess}}(H_{4\pi\alpha\delta}) = [0, +\infty).$

Let us look for eigenvalues of $H_{4\pi\alpha\delta}$ corresponding to radially symmetric eigenfunctions. That is, we consider the equation $H_{4\pi\alpha\delta}\psi = k^4\psi$ with the same convention (5.2) as above and look for special solutions of the form $\psi(x) = g(|x|)$ with $g \in H^2((0, \infty), r^2 dr)$. These eigenvalues are determined by eigenvalues of the one dimensional operator T_{α} in $L^2((0, \infty), r^2 dr)$ associated with the quadratic form

$$t_{\alpha}[g] := \int_{0}^{\infty} |r^{-1}[rg'(r)]'|^2 r^2 dr + \alpha |g(0)|^2, \quad \mathsf{D}(t_{\alpha}) := H^2((0,\infty), r^2 dr).$$

Using the unitary transform $U : L^2((0, \infty), r^2 dr) \to L^2((0, \infty), dr)$ which acts as (Ug)(r) := rg(r), the operator T_α is unitarily equivalent to the operator $\widetilde{T}_\alpha := UT_\alpha U^{-1}$ in $L^2((0, \infty))$ associated with the quadratic form

$$\tilde{t}_{\alpha}[f] := \int_{0}^{\infty} |f''(r)|^{2} dr + \alpha |f'(0)|^{2}, \quad \mathsf{D}(\tilde{t}_{\alpha}) := H^{2}((0,\infty)) \cap H^{1}_{0}((0,\infty))$$

One has

$$\begin{split} \widetilde{T}_{\alpha}f(r) &= f''''(r), \quad x \in (0,\infty), \\ \mathsf{D}\big(\widetilde{T}_{\alpha}\big) &= \big\{\eta \in H^4((0,\infty)) \cap H^1_0((0,\infty)) \ : \ f''(0) - \alpha f'(0) = 0\big\} \end{split}$$

The boundary values at 0 are well defined due to the Sobolev embedding $H^4((0, \infty)) \hookrightarrow C^3([0, \infty))$. Considering the general solution (5.3) of the differential equation $f'''' = k^4 f$ and imposing the integrability requirement (i.e. $C_1, C_4 = 0$) together with the boundary conditions at 0, we arrive at the condition

$$k = \frac{1}{2^{1/2}} \, \alpha \, e^{i\pi/4} \, .$$

In order to make this equation compatible with (5.2), we see that the point spectrum of T_{α} is not empty if, and only if, arg $\alpha \in (3\pi/4, 5\pi/4)$, or equivalently, $\Re \alpha < -|\Im \alpha|$. If this condition is satisfied, T_{α} possesses one eigenvalue corresponding to the right-hand side put to the power 4. Summing up,

$$\sigma_{\mathbf{p}}(T_{\alpha}) = \begin{cases} \left\{ -\frac{1}{4} \alpha^{4} \right\} & \text{if} \quad \Re \alpha < -|\Im \alpha|, \\ \\ \emptyset & \text{if} \quad \Re \alpha \ge -|\Im \alpha|. \end{cases}$$

Notice that the arguments of α for which the point spectrum is empty correspond to real points intersecting the half-axis $[0, +\infty) = \sigma_{ess}(T_{\alpha})$. So any eigenvalue of T_{α} , if it exists, is necessarily discrete.

In particular, choosing $\alpha := e^{i\theta}$ and varying $\theta \in (3\pi/4, 5\pi/4)$, any boundary point of the closed disk $\{\lambda \in \mathbb{C} : |\lambda| \le \frac{1}{4}\}$ except for the point $\lambda = \frac{1}{4}$ will be an eigenvalue of $H_{4\pi\alpha\delta}$. This proves the optimality of Theorem 1.1 for d = 3 because $\|4\pi\alpha\delta\|_{L^1(\mathbb{R}^3)} = 4\pi|\alpha| = 4\pi$.

Remark 5.2. One can also justify the optimality of the constant in the eigenvalue bound corresponding to the inclusion (1.3) for d = 3 in the limit $\varepsilon \to 0^+$ by considering the following family of regular but singularly scaled potentials

$$\delta_{\varepsilon}(x) := \begin{cases} \frac{1}{4\pi\varepsilon^3} & \text{if } |x| < \varepsilon, \\ 0 & \text{otherwise.} \end{cases}$$

Of course $\delta_{\varepsilon} \to \delta$ in the sense of distributions as $\varepsilon \to 0^+$. What is more, restricting to radially symmetric eigenfunctions, the point spectrum of $H_{4\pi\alpha\delta_{\varepsilon}}$ converges to the point spectrum of $H_{4\pi\alpha\delta}$ as $\varepsilon \to 0^+$. This can be checked by explicitly solving the radial parts of the differential equations of the eigenvalue problem in the separate intervals where δ_{ε} is constant and matching the solutions at the interface point $r = \varepsilon$.

ACKNOWLEDGMENT

The research was partially supported by the GACR grants No. 20-17749X.

REFERENCES

- [1] A. A. Abramov, A. Aslanyan, and E. B. Davies, Bounds on complex eigenvalues and resonances, J. Phys. A: Math. Gen. 34 (2001), 57-72.
- [2] M. S. Abramowitz and I. A. Stegun, eds., Handbook of mathematical functions, Dover, New York, 1965.
- [3] R. A. Adams, Sobolev spaces, Academic Press, New York, 1975.
- [4] J. Arazy and L. Zelenko, Virtual eigenvalues of the high order Schrödinger operator. I, Integral Equations Operator Theory 55 (2006), no. 2, 189–231.
- [5] J. Arazy and L. Zelenko, Virtual eigenvalues of the high order Schrödinger operator. II, Integral Equations Operator Theory 55 (2006), no. 3, 305–345.
- [6] J. M. Arrieta, F. Ferraresso, and P. D. Lamberti, Spectral analysis of the biharmonic operator subject to Neumann boundary conditions on dumbbell domains, Integral Equations Operator Theory 89 (2017), no. 3, 377–408.

HEMATISCHE | 1349

- [7] J.-C. Cuenin, Eigenvalue bounds for Dirac and fractional Schrödinger operators with complex potentials, J. Funct. Anal. 272 (2017), no. 7, 2987–3018.
- [8] D. E. Edmunds and W. D. Evans, Spectral theory and differential operators, Oxford University Press, Oxford, 1987.
- [9] T. Ekholm and A. Enblom, Critical Hardy-Lieb-Thirring inequalities for fourth-order operators in low dimensions, Lett. Math. Phys. 94 (2010), no. 3, 293–312.
- [10] A. Enblom, Estimates for eigenvalues of Schrödinger operators with complex-valued potentials, Lett. Math. Phys. 106 (2016), no. 2, 197–220.
- [11] W. D. Evans and R. T. Lewis, Counting eigenvalues of biharmonic operators with magnetic fields, Math. Nachr. 278 (2005), no. 12–13, 1524–1537.
- [12] L. Fanelli and D. Krejčiřík, Location of eigenvalues of three-dimensional non-self-adjoint Dirac operators, Lett. Math. Phys. 109 (2019), 1473–1485.
- [13] L. Fanelli, D. Krejčiřík, and L. Vega, Spectral stability of Schrödinger operators with subordinated complex potentials, J. Spectr. Theory 8 (2018), 575–604.
- [14] C. Förster and J. Östensson, Lieb-Thirring inequalities for higher order differential operators, Math. Nachr. 281 (2008), no. 2, 199–213.
- [15] R. L. Frank, Eigenvalue bounds for Schrödinger operators with complex potentials, Bull. Lond. Math. Soc. 43 (2011), 745–750.
- [16] I. S. Gradshteyn and I. M. Ryzhik, Table of integrals, series, and products, seventh edn., pp. xlviii+1171, Elsevier/Academic Press, Amsterdam, 2007.
- [17] A. Hulko, On the number of eigenvalues of the biharmonic operator on \mathbb{R}^3 perturbed by a complex potential, Rep. Math. Phys. 81 (2018), no. 3, 373–383.
- [18] T. Kato, Perturbation theory for linear operators, Springer-Verlag, Berlin, 1966.
- [19] T. Kato, Wave operators and similarity for some non-selfadjoint operators, Math. Ann. 162 (1966), 258–279.
- [20] J.-C. Kiik, P. Kurasov, and M. Usman, On vertex conditions for elastic systems, Phys. Lett. A 379 (2015), no. 34–35, 1871–1876.
- [21] E. H. Lieb and M. Loss, Analysis, Amer. Math. Soc., Providence, RI, 1997.
- [22] Y. Netrusov and T. Weidl, On Lieb–Thirring inequalities for higher order operators with critical and subcritical powers, Comm. Math. Phys. **182** (1996), no. 2, 355–370.
- [23] M. Reed and B. Simon, Methods of modern mathematical physics, II: Fourier analysis, self-adjointness, Academic Press, New York, 1975.
- [24] B. Simon, Quantum mechanics for Hamiltonians defined by quadratic forms, Princeton Univ. Press., New Jersey, 1971.
- [25] E. M. Stein, Interpolation of linear operators, Trans. Amer. Math. Soc. 83 (1956), 482-492.
- [26] T. Weidl, Remarks on virtual bound states for semi-bounded operators, Comm. Partial Differential Equations 24 (1999), no. 1–2, 25–60.

How to cite this article: Ibrogimov OO, Krejčiřík D, Laptev A. *Sharp bounds for eigenvalues of biharmonic operators with complex potentials in low dimensions*. Mathematische Nachrichten. 2021;**294**:1333–1349. https://doi.org/10.1002/mana.202000196