m3hsoln4.tex

M3H SOLUTIONS 4. 12.2.2016

Q1. (i) Pythagoras' theorem in ΔOMP gives

$$r^2 = u^2 + (\frac{1}{2}s)^2. \tag{a}$$

Then with v := r - u, Pythagoras in ΔMQR gives

$$w^{2} := v^{2} + (\frac{1}{2}s)^{2}$$

= $(r-u)^{2} + (\frac{1}{2}s)^{2} = r^{2} - 2ru + u^{2} + (\frac{1}{2}s)^{2}$
= $2r^{2} - 2ru = 2r(r-u) = 2rv$,

by (a).

(ii) The *n*-gon approximation to the circumference 2π of the unit circle is the perimeter $P_n = ns$; the 2*n*-gon approximation is $P_{2n} = 2nw$. Starting with n = 4, the unit square with $s = \sqrt{2}$, 6 iterations give

$$P = 6.283153355$$
 $(n = 256).$

The actual value of 2π correct to 9 places is

6.283185308.

The approximations increase with n, but one further iteration (on my calculator, a Casio fx-100) gives

```
6.283252652,
```

which is too high, so this is the best we can do by this method.

Q2 Fibonacci numbers.

The Fibonacci sequence satisfies the difference equation $u_n - u_{n-1} - u_{n-2} = 0$. The associated *characteristic equation* is $\lambda^2 - \lambda - 1 = 0$, with roots $\frac{1}{2}(1 \pm \sqrt{5})$. So the general solution is $u_n = c_1(\frac{1}{2}(1 + \sqrt{5}))^n + c_2(\frac{1}{2}(1 - \sqrt{5}))^n$. We can find c_1, c_2 from the initial conditions $u_0 = u_1 = 1$, giving

$$u_n = \frac{1}{2}(1 - 1/\sqrt{5})(\frac{1}{2}(1 - \sqrt{5}))^n + \frac{1}{2}(1 + 1/\sqrt{5})(\frac{1}{2}(1 + \sqrt{5}))^n.$$

For large n the second term dominates, and the result follows on division.

Q3 Long division: Fibonacci.

(i) If x is rational, x = m/n say:

(a) take off its integer part – so reducing to $0 \le m < n$,

(b) cancel m/n down to its lowest terms.

(ii) Now find the decimal expansion of m/n by the Long Division Algorithm. Let the remainders obtained by r_1, r_2, \ldots . The expansion *terminates* if some $r_k = 0$. It *recurs* if some remainder has *already occurred*. As there are only n-1 different possible non-zero remainders, the expansion must terminate (with remainder 0) or recur (with a remainder the first repeat of one of $1, 2, \ldots, n-1$) after at most n-1 places.

(ii) If x is a terminating decimal, x is a rational of the form $n + m/10^k$.

If x is a recurring decimal, say

$$x = n.a_1 \dots a_k b_1 \dots b_\ell \dots b_1 \dots b_\ell \dots,$$

x is $n.a_1...a_k$ (rational, above) +y, where writing

$$b := b_1/10 + \dots b_\ell/10^\ell$$

(rational, above), y is a geometric series with first term $b/10^k$ and common ratio $10^{-\ell}$. So

$$y = b.10^{-k} / (1 - 10^{-\ell}),$$

rational, so x is rational.

Combining with (i): x is rational iff its decimal expansion terminates or recurs.

(iii)

1/7 = 0.142857 recurring; 2/7 = 0.285714 rec.; 3/7 = 0.428571 rec.;

4/7 = 0.571428 rec.; 5/7 = 0.714285 rec.; 6/7 = 0.857142 rec.

These are the same six digits in each case, in cyclic order. What we see here is some connection between the denominator 7 and the base 10. For theoretical background here, see e.g. G. H. Hardy & E. M. Wright, An introduction to the theory of numbers, 6th ed., OUP, 2008 (1st ed. 1938),

Ch. IX: The representation of numbers by decimals.

But note that from the point of view of Number Theory, the only natural way to expand a real number is as a *continued fraction*: see e.g. Hardy & Wright, Ch. X: Continued fractions.

NHB