

**M3PM16/M4PM16 SOLUTIONS 8. 22.3.2012**

Q1. The numbers  $1, 2, \dots, n$  include:  $[n/p]$  multiples of  $p$ ;  $[n/p^2]$  multiples of  $p^2$ ; etc. Multiplying  $1, 2, \dots, n$  to get  $n!$ , the prime powers add, as required.

Q2. This follows from  $N = (2n)!/(n!)^2$  and Q1.

Q3. If  $[2x] = 2n+1$  is odd,  $2n+1 \leq 2x < 2n+2$ ,  $n + \frac{1}{2} \leq x < n+1$ ,  $[x] = 2n$ , so  $[2x] - 2[x] = 2n+1 - 2n = 1$ . If  $[2x] = 2n$  is even,  $2n \leq 2x < 2n+1$ ,  $n \leq x < n + \frac{1}{2}$ ,  $[2x] - 2[x] = 2n - 2n = 0$ .

In Q2, each term with  $p^m > 2n$  is 0. There are  $[\log 2n/\log p]$  other terms, each 0 or 1 by above, so  $k(p) \leq [\log 2n/\log p] \leq \log 2n/\log p$ .

Q4 (*Bertrand's postulate: Erdős' proof of 1932*; see HW §22.3).

We can check this by hand for  $n \leq 2^9 = 512$  (below). For  $n > 2^9$ , we assume not and derive a contradiction. So, we assume there is no prime  $p$  with  $n < p \leq 2n$ .

Write  $N := \binom{2n}{n}$ . Then if  $p$  is a prime divisor of  $N$ ,  $p \leq 2n$ , so  $p \leq n$  by hypothesis. Also  $k(p) \geq 1$  in Q2.

Assume  $\frac{2}{3}n < p \leq n$  (we shall see that this case cannot occur). Then  $2p \leq 2n < 3p$ ,  $p^2 > \frac{4}{9}n^2 > n \cdot \frac{4}{9} \cdot 512 > 2n$ :  $2n/p^2 < 1$ , so  $[2n/p^2] = 0$ , so also  $[2n/p^m] = 0$  for  $m \geq 2$ . So

$$k(p) := \sum_m ([2n/p^m] - 2[n/p^m]) = [2n/p] - 2[n/p].$$

As  $2 \leq 2n/p < 3$  (above),  $[2n/p] = 2$ . As  $1 \leq n/p < 3/2$  (above),  $[n/p] = 2$ . Combining,  $k(p) = 0$ . But  $k(p) \geq 1$  (above), so no such  $p$  exists.

We now have  $p \leq \frac{2}{3}n$  for every prime factor  $p$  of  $N$ . So

$$\sum_{p|N} \log p \leq \sum_{p \leq \frac{2}{3}n} \log p = \theta\left(\frac{2}{3}n\right) \leq \frac{4}{3} \log 2n, \quad (1)$$

by Chebyshev's Upper Estimate (III.2).

If  $k(p) \geq 2$ ,  $2 \log p \leq k(p) \log p \leq \log(2n)$  by Q3. So

$$\log p \leq \log \sqrt{2n} : \quad p \leq \sqrt{2n}.$$

So there are at most  $\sqrt{2n}$  such  $p$ . So

$$\sum_{k(p) \geq 2} k(p) \log p \leq \sqrt{2n} \log(2n)$$

(at most  $\sqrt{2n}$  terms; in each,  $p \leq 2n$ , as  $p|N = \binom{2n}{n}$ ). So by (1),

$$\begin{aligned} \log N &= \sum_{p \leq 2n} k(p) \log p = \sum_{k(p)=1} \log p + \sum_{k(p) \geq 2} k(p) \log p \\ &\leq \sum_{p|N} \log p + \sqrt{2n} \log(2n) \leq \frac{4}{3} \log 2n + \sqrt{2n} \log(2n). \end{aligned} \quad (2)$$

Also,  $N$  is the largest term in the binomial expansion of  $(1+1)^{2n} = 2^{2n}$ , so  $2^{2n} = 2 + \binom{2n}{1} + \dots + \binom{2n}{2n-1} \leq 2nN$ . So by (2)  $2n \log 2 \leq \log(2n) + \log N \leq \frac{4}{3} \log 2n + (1 + \sqrt{2n}) \log(2n)$ , giving

$$2n \log 2 \leq 3(1 + \sqrt{2n}) \log(2n). \quad (3)$$

Write

$$x := \frac{\log(n/512)}{10 \log 2} > 0$$

(as  $n > 512$ ):  $10x \log 2 = \log(n/2^9)$ ,

$$10(1+x) \log 2 = \log(n/2^9) + \log(2^{10}) = \log(2n) : \quad 2n = 2^{10(1+x)}.$$

So (3) is  $2^{10(1+x)} \log 2 \leq 3(1 + 2^{5(1+x)}) \cdot 10(1+x) \log 2$ :  
 $2^{10(1+x)} \leq 30(1 + 2^{5+5x})(1+x)$ . Divide by  $2^{10} \cdot 2^{5x}$ :

$$\begin{aligned} 2^{5x} &\leq 30 \cdot 2^{-5} (1 + 2^{-5-5x})(1+x) \leq 30 \cdot 2^{-5} (1 + 2^{-5})(1+x) \quad (x > 0) \\ &< (1 - 2^{-5})(1 + 2^{-5})(1+x) \quad (30 \cdot 2^{-5} < 31 \cdot 2^{-5} = 1 - 2^{-5}) \\ &< 1 + x. \end{aligned} \quad (4)$$

But  $2^{5x} = \exp(5x \log 2) > 1 + 5x \log 2$  ( $e^x > 1 + x$  for  $x > 0$ )  $> 1 + x$  ( $5 \log 2 = \log 2^5 > \log e = 1$ ). But this contradicts (4). So for each  $n > 512$  there is indeed a prime  $p$  with  $n < p \leq 2n$ .

For the early primes: each of the primes

$$2, 3, 5, 7, 13, 23, 43, 83, 163, 317, 631$$

is less than twice its predecessor. So for any  $n \leq 630$ , at least one such  $p$  satisfies  $n < p \leq 2n$ . Combining gives the result.

NHB