## m3pm16l9.tex Lecture 9. 31.1.2013

**Lemma**. If  $A(x) := \sum_{n \leq x} a_n$ ,  $B(x) := \sum_{n \leq x} b_n$ ,

$$\sum_{n \le x} (a * b)(n) = \sum_{jk \le x} a_j b_k = \sum_{j \le x} a_j B(x/j) = \sum_{k \le x} b_k A(x/k).$$

Proof.

$$\sum_{n \le x} (a * b)(n) = \sum_{n \le x} \sum_{jk=n} a_j b_k = \sum_{jk \le x} a_j b_k = \sum_{j \le x} a_j \sum_{k \le x/j} b_k = \sum_{j \le x} a_j B(x/j),$$

and symmetrically. //

Defn. Call a multiplicative if a(.) is not  $\equiv 0$  and

$$a(mn) = a(m)a(n) \qquad \text{for } (m,n) = 1$$

((m,n) = gcd of m and n: (m,n) = 1 means m, n are coprime - have no common factors.

Call a completely multiplicative if it is not  $\equiv 0$  and

$$a(mn) = a(m)a(n) \quad \forall \ m, n.$$

**Propn.** (i) If a is multiplicative, a(1) = 1. (ii) If a, b are multiplicative, so is a \* b.

*Proof.* (i) As (n, 1) = 1 for all n, a(n)a(1) = a(n). As a is not  $\equiv 0$ ,  $a(n) \neq 0$  for some n. Then cancelling gives a(1) = 1.

(ii) Take m, n with (m, n) = 1. As m, n have no common factors, every divisor r of mn is uniquely expressible as r = jk with j|m and k|n. Then also j, k have no common factors, so (j, k) = 1. Similarly, (m/j, n/k) = 1. So

$$(a*b)(n) = \sum_{r|mn} a(r)b(mn/r) = \sum_{j|m} \sum_{k|n} a(jk)b(\frac{m}{j}, \frac{n}{k}) = \sum_{j|m} \sum_{k|n} a(j)a(k)b(\frac{m}{j})b(\frac{n}{k})$$

(as both a and b are multiplicative)

$$= (a * b)(m)(a * b)(n).$$
 //

**Cor.** If f is multiplicative, so is F := f \* u:  $F(n) = \sum_{d|n} f(d)$ .

There is a converse: if  $F(n) = \sum_{d|n} f(d)$  is multiplicative, so is f (Problems 5, Q3).

## §4. Euler Products

Throughout, write p for a prime, P for the set of primes,

**Theorem (Euler).** If a is completely multiplicative with  $\sum_{1}^{\infty} |a_n| < \infty$ , then (i)  $\sum_{n=0}^{\infty} a_n \neq 0$ :

(i) 
$$\sum_{1}^{\infty} a_n \neq 0$$
,  
(ii)  $\sum_{1}^{\infty} a_n = \prod_p 1/(1 - a_p)$ .

*Proof.* By I.5,  $\prod_p (1 - a_p)$  converges to a non-zero value (as  $\sum |a_n| < \infty$ ); thus so does  $\prod_p 1/(1 - a_p)$ .

Fix N; write P[N] for the set of primes  $p \leq N$ ,  $E_N$  for the set of integers with all prime factors in P[N],  $E_N^*$  for the remaining natural numbers,

$$T_N := \prod_{p \in P[N]} 1/(1 - a_p) = \prod_{p \in P[N]} (1 + a_p + a_p^2 + \ldots).$$

Multiply out. Each  $n \in E_N$  appears on RHS exactly once, by FTA (I.1). So

$$T_N = \sum_{n \in E_N} a_n$$

As  $\{1, 2, ..., N\} \subset E_N, E_N^* \subset \{N + 1, N + 2, ...\}$ , so

$$|\sum_{1}^{\infty} a_n - T_N| = |\sum_{n \in E_N^*} a_n| \le \sum_{n > N} |a_n| \to 0 \qquad (N \to \infty). \qquad //$$

The special case  $a_n \equiv 1/n^s$  gives

**Theorem (Euler).**  $\zeta(s) = \prod_p 1/(1 - 1/p^s) \ (Res > 1).$ *Proof.* 

$$RHS = \prod_{p} (1 + p^{-s} + p^{-2s} + \dots) = \sum_{k, p_1, \dots, p_k} p_1^{-n_1 s} p_2^{-n_2 s} \dots p_k^{-n_k s} = \sum_n n^{-s} = \zeta(s)$$

by FTA, as each  $n=p_1^{n_1}\dots p_k^{n_k},$  uniquely. //