m3pm16l30.tex Lecture 31. 25.3.2014.

Recall (IV.2 L27) the complex Stirling formula for Γ , giving for its logarithmic derivative $\Gamma'(z)/\Gamma(z) = O(\log z)$ in any region avoiding 0 and the negative half-line. So in the critical strip,

$$\Gamma'(s)/\Gamma(s) = O(\log t).$$

By the partial fraction expansion (IV.2 L27)

$$-\frac{\zeta'(s)}{\zeta(s)} = -B + \frac{1}{s-1} - \frac{1}{2}\log\pi + \frac{1}{2}\frac{\Gamma'(\frac{1}{2}s+1)}{\Gamma(\frac{1}{2}s+1)} - \sum_{\rho} \left(\frac{1}{s-\rho} + \frac{1}{\rho}\right).$$

By (*) (L30), the real part of the sum here is ≥ 0 in (ZFR). So we can *discard* it here when estimating the *real part* of the LHS above. Combining:

$$-Re\zeta'(s)/\zeta(s) \le O(\log t) \qquad (t \ge 4, \sigma \ge 1 - 4c/\log t). \qquad (**)$$

Note. Here " \leq " is crucial (we need a *one-sided* bound to use Borel-Carathéodory). We do *not* mean "<<".

In what follows (and indeed, throughout), we may take t arbitrarily large. Fix $s = \sigma + it$ with $t \ge 5$, $\sigma \ge 1 - c/\log t$ (to keep within the zero-free region). Write

$$\eta := c/\log t, \qquad s_0 := 1 + \eta + it$$

(so η is small). For w in the disc $|w| \leq 4\eta$ ($w = u + iv = re^{i\theta}$, $|r| \leq 4\eta$ small)

$$\sigma' + it' := s_0 + w = 1 + \eta + u + it + iv$$

satisfies

$$t' \ge 4$$
 $(t' = t + v, t \ge 5, v \text{ small}),$
 $\sigma' \ge 1 - \frac{4c}{\log t'}$

 $(\sigma' = 1 + \eta + u = 1 + c/\log t + u; u \text{ small}, t' - t + v, v \text{ small})$. So by (**),

$$-Re \zeta'(s_0+w)/\zeta(s_0+w) \le 2K\log t,$$

for some K.

Now write

$$F(w) := -\frac{\zeta'(s_0 + w)}{\zeta(s_0 + w)} + \frac{\zeta'(s_0)}{\zeta(s_0)}$$

This satisfies the hypotheses of the Borel-Carathódory theorem (IV.2 L29) in the disc $|w| \leq 4\eta$, with

$$A := 2K \log \tau + |\zeta'(s_0)/\zeta(s_0)|.$$

But

$$|s - s_0| \le 2\eta$$
, $(s - s_0 = \sigma - 1 \ge -2\eta$, and $s - s_0 \le 0$).

By the Borel-Carathéodory theorem, with $R = 4\eta$, $r = 2\eta$:

$$\left|\frac{\zeta'(s)}{\zeta(s)}\right| \le 4K\log t + 3\left|\frac{\zeta'(s_0)}{\zeta(s_0)}\right| \qquad \left(\frac{2r}{R-r} = 2, \frac{R=r}{R-r} = 3\right).$$

But as $s_0 = 1 + \eta + i\tau$,

$$\left|\frac{\zeta'(s_0)}{\zeta(s_0)}\right| = \left|\sum_{1}^{\infty} \Lambda(n)/n^{1+\eta+i\tau}\right| \le \sum_{1}^{\infty} \Lambda(n)/n^{1+\eta}$$
$$= 1/\eta + O(1)$$

(as $-\zeta'/\zeta$ has a simple pole at 1 with residue 1)

$$<<\log t$$
 $(\eta := c/\log t).$ //

5. PNT with remainder. We follow Montgomery and Vaughan [MV], 6.2.

Theorem (PNT with Remainder). For some c > 0,

$$\begin{split} \psi(x) &= x + O(x \exp\{-c\sqrt{\log x}\}, \quad \theta(x) = x + O(x \exp\{-c\sqrt{\log x}\}, \\ \pi(x) &= li(x) + O(x \exp\{-c\sqrt{\log x}\}. \end{split}$$

Proof. The equivalence follows as in the Equivalence Theorem for PNT (III.2 L17). To prove the first: by 'Perron for $-\zeta'/\zeta$ ' (Th. 3, IV.1 L27),

$$\psi(x) = \frac{1}{2\pi i} \int_{k-iT}^{k+iT} -\frac{\zeta'(s)}{\zeta(s)} \frac{x^s}{s} ds + O\left(\log x \left[1 + \frac{x\log T}{T}\right]\right)$$

for $k := 1 + 1/\log x$. By (ZFR) (IV.3 L29-30), for some $c_0 > 0$ s = 1 is the only singularity of the integrand in the rectangle

$$|\tau| \le T$$
, $1 - c_0 / \log T \le \sigma \le k = 1 + 1 / \log x$.