

## §3. Holomorphy

**Theorem.** A Dirichlet series is holomorphic within its half-plane of convergence, with derivative given by termwise differentiation. If  $F(s) := \sum_1^\infty a_n/n^s$  for  $\sigma > \sigma_c$ , then  $F'(s) = -\sum_1^\infty \log n \, a_n/n^s$ .

*Proof.* Choose  $\alpha > \sigma_c$  and write  $b_n := a_n/n^\alpha$ . As above,  $b_n$  is bounded (by  $M$ , say). Write  $F(s) = G(s - \alpha)$ ,  $G(s) := \sum_1^\infty b_n/n^s$ . Write  $G_N(s) := \sum_1^N b_n/n^s = \sum_1^N b_n e^{-s \log n}$ . Then  $G'_N(s) = -\sum_1^N \log n \, b_n/n^s$ .

Take  $\delta > 0, R > 0, K > 0, \Gamma$  the rectangle with sides  $\sigma = \delta, \sigma = K, t = \pm R, E$  its interior. By II.1, (\*\*),

$$|G(s) - G_N(s)| \leq \frac{M}{N^\sigma} \left( \frac{|s|}{\sigma} + 1 \right).$$

For  $s \in E$ ,

$$\frac{|s|}{\sigma} \leq \frac{\sigma + |t|}{\sigma} = 1 + \frac{|t|}{\sigma} \leq 1 + \frac{R}{\sigma}.$$

So

$$|G(s) - G_N(s)| \leq \frac{M}{N^\delta} \left( 2 + \frac{R}{\delta} \right) \rightarrow 0 \quad (N \rightarrow \infty),$$

uniformly on  $\Gamma \cup E$ , which is compact. As each  $G_N$  is holomorphic by I.2,  $G$  is holomorphic. As each  $s$  with  $\sigma > 0$  is in some  $E$ ,  $G$  is holomorphic on  $\sigma > 0$ , so  $F$  is holomorphic on  $\sigma > \alpha$ . Then  $G'_N \rightarrow G'$  by I.2, so as  $D(n^{-s}) = D(e^{-s \log n}) = -\log n \, n^{-s}$ ,  $F'(s) = -\sum_1^\infty \log n \, a_n/n^s$ . Similarly for Dirichlet integrals: if  $I_X(s) := \int_1^X f(x) dx/x^{1+s}$ , then  $I'_X(s) = -\int_1^X f(x) \log x dx/x^{1+s}$  by differentiating under the integral sign. //

*Example.*

$$\zeta(s) = \sum_1^\infty \frac{1}{n^s}, \quad \zeta'(s) = -\sum_1^\infty \log n/n^s \quad (\sigma > 1).$$

By integrating by parts,

$$\int_1^\infty \frac{\log x}{x^\sigma} dx = \frac{1}{(\sigma - 1)^2} \quad (\sigma > 1).$$

Hence as in I.4,

$$-\zeta'(\sigma) \leq \frac{1}{(\sigma-1)^2}.$$

#### §4. Convolutions

Absolutely convergent series may be rearranged. So if

$$F_a(s) := \sum_1^\infty a_n/n^s, \quad F_b(s) := \sum_1^\infty b_n/n^s,$$

then in the half-plane where both converge absolutely

$$F_a(s)F_b(s) = \left(\sum_{i=1}^\infty \frac{a_i}{i^s}\right)\left(\sum_{j=1}^\infty \frac{b_j}{j^s}\right) = \sum_{ij} \frac{a_i b_j}{i^s j^s} = \sum_{n=1}^\infty \frac{c_n}{n^s},$$

where

$$c_n := \sum_{ij=n} a_i b_j = \sum_{i|n} a(i)b(n/i).$$

The series  $c = (c_n)$  so obtained is called the *Dirichlet convolution* of  $a$  and  $b$ :

$$c = a * b.$$

Write  $e_i := (\delta_{1n})$  (the Kronecker delta: 1 if  $n = 1$ , 0 otherwise). Then  $a * e_1 = a$ :  $e_1$  acts as an identity.

Dirichlet convolutions have the properties:

$a * b = b * a$  – commutativity;

$a * (b + c) = a * b + a * c$  – distributivity;

$a * (b * c) = (a * b) * c$  – associativity.

Note also:  $u := (u_n)$ , where  $u_n := 1$  for all  $n$ , so  $u$  has Dirichlet series

$$\zeta(s) := \sum_1^\infty 1/n^s; \quad (u, \zeta)$$

$d := (d_n)$ , the *divisor function*, where  $d_n := \sum_{d|n} 1$  is the number of divisors of  $n$ . Then

$$(u * u)_n = \sum_{d|n} u(d)u(n/d) = \sum_{d|n} 1 = d(n) : \quad u * u = d.$$

So one has the important Dirichlet series

$$\zeta(s)^2 = \sum_{n=1}^\infty d_n/n^s. \quad (d, \zeta^2)$$