

### III. THE PRIME NUMBER THEOREM AND ITS RELATIVES

#### §1. The Prime Number Theorem (PNT)

PNT states that

$$\pi(x) := \sum_{p \leq x} 1 \quad \sum li(x) := \int_2^x dt / \log t \sim x / \log x \quad (x \rightarrow \infty) \quad (PNT)$$

This was conjectured on numerical grounds by GAUSS (c. 1799; letter of 1848) and A. M. LEGENDRE (1752-1833; in 1798, *Essai sur la Théorie des Nombres*).

In 1737 L. EULER (1707-1783) found his Euler product, linking the primes to  $\sum_{n=1}^{\infty} 1/n^{\sigma}$  for real  $\sigma$  (later the Riemann zeta function).

In 1850-51, P. L. CHEBYSHEV (= TCHEBYSHEV, etc., 1821-1894) made two great strides (III.2):

- (i)  $\pi(x) \asymp x / \log x$ , i.e.  $C_1 x / \log x \leq \pi(x) \leq C_2 x / \log x$  for some  $0 < C_1 \leq C_2 < \infty$  and all  $x \geq X$ ;
- (ii)

$$\liminf \pi(x) / \frac{x}{\log x} \leq 1 \leq \limsup \pi(x) / \frac{x}{\log x}$$

– so if the limit exists (which we shall prove!), it must be 1.

In 1859 B. RIEMANN (1826-66) studied

$$\zeta(s) := \sum_{n=1}^{\infty} 1/n^s \quad (s \in \mathbb{C})$$

using Complex Analysis (M2PM3), then still fairly new, developed by A. L. CAUCHY (1789-1857), 1825-29. He showed the critical relevance of the *zeros* of  $\zeta(s)$  to the *distribution of primes*. We shall show that:

- (i)  $\zeta$  can be continued analytically from  $\operatorname{Re} s > 1$  to the whole complex plane  $\mathbb{C}$ , where it is holomorphic except for a simple pole at 1 of residue 1 (III.3);
- (ii) The only zeros of  $\zeta$  outside the *critical strip*

$$0 < \sigma = \operatorname{Re} s < 1$$

are the so-called *trivial zeros*  $-2, -4, \dots, -2n, \dots$  (trivial in that they follow from the *functional equation* for  $\zeta$  – see III.7);

- (iii) PNT is closely linked to non-vanishing of  $\zeta$  on the 1-line (III.4):

$$\zeta(1 + it) \neq 0.$$

Indeed, PNT is equivalent to this.

The *Riemann Hypothesis* (RH) of 1859 is that the only zeros of  $\zeta$  in the critical strip are on the *critical line*

$$\sigma = \frac{1}{2}. \quad (RH)$$

RH is still open, and is the most famous and important open question in Mathematics. Its resolution would have vast consequences for prime-number theory (especially error terms in PNT – see e.g. J Ch. 5). It is so hard that proving theorems *conditional on RH* (i.e., assuming it is true) is respectable in Analytic Number Theory.

PNT was proved independently in 1896 by J. HADAMARD (1865-1963, French) and Ch. de la Vallée Poussin (1866-1962, Belgian). Both used Complex Analysis and  $\zeta$ .

Since counting primes relates to  $\mathbb{N} (\subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C})$ , it seemed strange and unaesthetic to use complex methods. Great efforts were made to provide an *elementary proof*, over half a century.

Elementary proofs of PNT were found in 1948 by Paul ERDÖS (1913-1996, Hungarian) and Atle SELBERG (1917-2009, Norwegian). There is a full account by J. Spencer and R. Graham in *The Mathematical Intelligencer*, **31.3** (2009), 18-23. Erdős gave an elementary proof of a result of Chebyshev (proof of Bertrand's postulate: Problems 8). Selberg told him the day after seeing it that he could use it to complete an elementary proof of PNT. Erdős proposed collaboration but Selberg declined; their papers were published separately in 1949.

Proofs of ANT by complex methods are in all the books on ANT, including J Ch. 3, A, R, D. Elementary proofs of PNT are harder; see e.g. HW Ch. XXII (22.14-16), J Ch. 6, A Ch. 4, R Ch. 13.

Error estimates in PNT are very important. Naturally, complex methods give better error estimates than elementary ones. Error estimates depend on *zero-free regions* of  $\zeta$  (to the left of the 1-line, in the critical strip) – the bigger, the better; see IV.3.