m3pm16soln1.tex

## M3PM16/M4PM16 SOLUTIONS 1. 22.1.2015

Q1 (HW, Th. 135). (i) If x is a terminating decimal, x is a rational of the form  $n + m/10^k$ . If x is a recurring decimal, say

 $x = n.a_1 \dots a_k b_1 \dots b_\ell \dots b_1 \dots b_\ell \dots,$ 

x is  $n.a_1...a_k$  (rational, above) +y, where writing

$$b := b_1/10 + \dots b_\ell/10^\ell$$

(rational, above), y is a GP with first term  $b/10^k$  and common ratio  $10^{-\ell}$ . So

$$y = b.10^{-k} / (1 - 10^{-\ell}),$$

rational, so x is rational.

If x is rational, x = m/n say:

(a) take off its integer part – so reducing to  $0 \le m < n$ ,

(b) cancel m/n down to its lowest terms.

(ii) Now find the decimal expansion of m/n by the Long Division Algorithm. Let the remainders obtained by  $r_1, r_2, \ldots$ . The expansion *terminates* if some  $r_k = 0$ . It *recurs* if some remainder has *already occurred*. As there are only n-1 different possible non-zero remainders, the expansion must terminate (with remainder 0) or recur (with a remainder the first repeat of one of  $1, 2, \ldots, n-1$ ) after at most n-1 places. (As 1/7 shows, all n-1 places may be needed.)

(iii) Similarly with 10 replaced by 2,3, ...

Q2 ([L], 12-13). For  $x \ge 4$ ,

$$li(x) := \int_2^x \frac{du}{\log u} = \left[\frac{u}{\log u}\right]_2^x - 2\int_2^x d\left(\frac{1}{\log u}\right) = \frac{x}{\log x} - \frac{2}{\log 2} + \int_2^x \frac{du}{\log^2 u}.$$

But

$$0 < \int_{2}^{x} \frac{du}{\log^{2} u} = \int_{2}^{\sqrt{x}} + \int_{\sqrt{x}}^{x} < \int_{2}^{\sqrt{x}} \frac{2}{\log^{2} 2} + \int_{\sqrt{x}}^{x} \frac{du}{\log^{2} x}$$
$$= \frac{\sqrt{x} - 2}{\log^{2} 2} + \frac{x - \sqrt{x}}{\frac{1}{4}\log^{2} x} < \frac{\sqrt{x}}{\log^{2} 2} + \frac{4x}{\log^{2} x} = o\left(\frac{x}{\log x}\right)$$

(as  $\log \sqrt{x} = \frac{1}{2} \log x$ ). Combining,

$$li(x) = \frac{x}{\log x} + o(\frac{x}{\log x}): \qquad li(x) / \frac{x}{\log x} \to 1.$$

Q3 ([L], 13-14). For  $x \ge 4$ , as before, integrating by parts m + 1 times,

$$\begin{split} li(x) &= \frac{x}{\log x} + \frac{1!x}{\log^2 x} + \ldots + \frac{m!x}{\log^{m+1}x} + const + (m+1)! \int_2^x \frac{du}{\log^{m+2}u} :\\ 0 &< \int_2^x \frac{du}{\log^{m+2}u} = \int_2^{\sqrt{x}} + \int_{\sqrt{x}}^x < \int_2^{\sqrt{x}} \frac{du}{\log^{m+2}2} + \int_{\sqrt{x}}^x \frac{du}{\log^{m+2}(\sqrt{x})} \\ &< \frac{\sqrt{x}-2}{\log^{m+2}2} + \frac{x-\sqrt{x}}{2^{-m-2}\log^{m+2}x} = o\Big(\frac{x}{\log^{m+1}x}\Big) :\\ li(x) - \Big(\frac{x}{\log x} + \frac{1!x}{\log^2 x} + \ldots + \frac{(m-1)!x}{\log^m x}\Big) = \frac{m!x}{\log^{m+1}x}(1+o(1)). \end{split} //$$

Note. Numerical evidence shows that li(x) gives a much better approximation than  $x/\log x$ , in line with Q1, Q2, and we shall prefer it – particularly in PNT with any error term – see Ch. IV and Problems 2.

Q4 ([L], 214-5). Taking  $x = p_n$  in  $\pi(x) := \sum_{p \le x} 1$  gives  $\pi(p_n) = \sum_{p \le p_n} 1 = n.$ 

By PNT,  $\pi(x) \sim x/\log x$ , so  $n \sim p_n/\log p_n$ :

$$\frac{n\log p_n}{p_n} \to 1. \tag{i}$$

Taking logs of (i),  $\log n + \log \log p_n - \log p_n \to 0$ . Dividing this by  $\log p_n$ ,

$$\frac{\log n}{\log p_n} + \frac{\log \log p_n}{\log p_n} - 1 \to 0.$$

But  $\log x = o(x)$ , so  $\log \log p_n = o(\log p_n)$ , so this says

$$\frac{\log n}{\log p_n} \to 1. \tag{ii}$$

Multiply (i) and (ii):  $n \log n / \log p_n \to 1$ , i.e.  $p_n \sim n \log n$ . // NHB