

# SOLUTIONS 10 12.12.2013

Q1. With  $\pi$  as in Q2,

$$\pi_i p_{i,i+1} = \frac{1}{\binom{2d}{d}} \binom{d}{i}^2 \cdot \left(\frac{d-i}{d}\right)^2 = \frac{1}{\binom{2d}{d}} \binom{d-1}{i}^2,$$

as in Problems 9 Q3, and similarly

$$\pi_{i+1} p_{i+1,i} = \frac{1}{\binom{2d}{d}} \binom{d}{i+1}^2 \cdot \left(\frac{i+1}{d}\right)^2 = \frac{1}{\binom{2d}{d}} \binom{d-1}{i}^2,$$

proving detailed balance, and so reversibility. Assuming reversibility, we can use detailed balance to calculate the invariant distribution, again using the calculation in Problems 9 Q2:

$$\pi_i = \frac{\pi_0}{\left(\frac{1}{d}\right)^2} \cdot \frac{\left(1 - \frac{1}{d}\right)^2}{\left(\frac{2}{d}\right)^2} \cdot \dots \cdot \frac{\left(1 - \frac{i-1}{d}\right)^2}{\left(\frac{i}{d}\right)^2} = \pi_0 \cdot \frac{(d(d-1) \dots (d-i+1))^2}{(1 \cdot 2 \dots i)^2} = \pi_0 \binom{d}{i}^2.$$

Then  $\sum_i \pi_i = 1$  gives

$$\pi_0 \sum_i \binom{d}{i}^2 = \pi_0 \binom{2d}{d}^2 = 1, \quad \pi_0 = 1 / \binom{2d}{d}, \quad \pi_i = \binom{d}{i}^2 / \binom{2d}{d}. \quad //$$

Q2.  $\pi_i = 1/\mu_i$  by the Erdős-Feller-Pollard theorem (L29), so

$$\mu_0 = 1/\pi_0 = \binom{2d}{d}.$$

By Stirling's formula,

$$\mu_0 \sim \frac{\sqrt{2\pi} e^{-2d} 2d^{2d+\frac{1}{2}}}{(\sqrt{2\pi} e^{-d} d^{d+\frac{1}{2}})^2} = \frac{4^d}{\sqrt{\pi d}}.$$

Now as  $d$  is already very large (of the order of Avogadro's number  $6 \times 10^{23}$ ),  $4^d$  is astronomically vast – effectively infinite.

The interpretation of this in Statistical Mechanics is that  $\mu_0$  is the mean recurrence time of state 0, when all the  $2d$  gas molecules are in one half of the container. Although this state is certain to recur (indeed, infinitely often), its mean recurrence time is so vast as to be effectively infinite – which explains why we do not see such states recurring in practice! This reconciles the theoretical reversibility of the model with the irreversible behaviour we observe when gases diffuse, etc. This was the Ehrenfests' motivation for their model, in 1912.

*Note.* Relevant here is the concept of *entropy* – a measure of disorder. This was introduced by Rudolf CLAUSIUS (1922-1888), in 1865, who formulated the First Law of Thermodynamics (Law of Conservation of Energy) and Second Law of Thermodynamics (entropy increases – things become more disordered):

1. Die Energie der Welt ist konstant (The energy of the world [the universe] is constant).
2. Die Entropie der Welt strebt einem Maximum zu (The entropy of the world [the universe] strives towards a maximum).

Q3. Branching processes.

(i)  $Z_2$  is the sum of a random number,  $Z_1$ , of independent copies of  $Z$ . So

$$P_2(s) := E[s^{Z_2}] = \sum_{k=0}^{\infty} E[s^{Z_2} | Z_1 = k] P(Z_1 = k).$$

Now when  $Z_1 = k$ ,  $Z_2$  is a sum of  $k$  independent copies of  $Z$ , each with PGF  $P(s)$ , so has (conditional) PGF  $P(s)^k$ . So

$$P_2(s) = \sum_0^{\infty} p_k P(s)^k = P(P(s)).$$

(ii) Similarly, or by induction on  $n$ ,  $Z_n$  has PGF  $P_n$ .

(iii)

$$P'_n(s) = P'(P_{n-1}(s)).P'_{n-1}(s).$$

So letting  $s = 1$  ( $R > 1$ ), or  $s \uparrow 1$  ( $R = 1$ ) and using Abel's Continuity Theorem, since  $P_{n-1}$ , being a PGF, has value 1 at 1,  $P'_n(1) = P'(1).P'_{n-1}(1) = \mu.P'_{n-1}(1)$ , so by induction

$$P'_n(1) = \mu^n : \quad E[Z_n] = \mu^n.$$

NHB