## ma414l17.tex Lecture 17. 10.3.2011

*Example.* We calculate  $\int W(u)dW(u)$ . We start by approximating the integrand by a sequence of simple functions.

$$X_n(u) = \begin{cases} W(0) = 0 & \text{if} & 0 \le u \le t/n, \\ W(t/n) & \text{if} & t/n < u \le 2t/n, \\ \vdots & \vdots & \\ W((n-1)t/n) & \text{if} & (n-1)t/n < u \le t. \end{cases}$$

By definition,

$$\int_0^t W(u) dW(u) = \lim_{n \to \infty} \sum_{k=0}^{n-1} W(kt/n) (W((k+1)t/n) - W(kt/n)).$$

Replacing W(kt/n) by  $\frac{1}{2}(W((k+1)t/n) + W(kt/n)) - \frac{1}{2}(W((k+1)t/n) - W(kt/n)))$ , the RHS is

$$\sum \frac{1}{2} (W((k+1)t/n) + W(kt/n)) \cdot (W((k+1)t/n) - W(kt/n)) - \sum \frac{1}{2} (W((k+1)t/n) - W(kt/n)) \cdot (W((k+1)t/n) - W(kt/n))) \cdot (W((k+1)t/n) - W(kt/n))) \cdot (W((k+1)t/n) - W(kt/n)) \cdot (W(k+1)t/n) - W($$

The first sum is  $\sum \frac{1}{2}(W((k+1)t/n)^2 - W(kt/n)^2)$ , which telescopes (as a sum of differences) to  $\frac{1}{2}W(t)^2$  (W(0) = 0). The second sum is

 $\frac{1}{2}\sum (W(k+1)t/n) - W(kt/n))^2$ , an approximation to the quadratic variation of W on [0, t], which tends to  $\frac{1}{2}t$  by Lévy's theorem on the QV. Combining,

$$\int_0^t W(u)dW(u) = \frac{1}{2}W(t)^2 - \frac{1}{2}t.$$

Note the contrast with ordinary (Newton-Leibniz) calculus! Itô calculus requires the second term on the right – the Itô correction term – which arises from the quadratic variation of W.

One can construct a closely analogous theory for stochastic integrals with the Brownian integrator W above replaced by a square-integrable martingale integrator M. The properties above hold, with the Lemma (i) replaced by

$$E[(\int_0^t X(u)dM(u))^2] = E[\int_0^t X(u)^2 d\langle M \rangle(u)].$$

The natural class of integrands X to use here is the class of predictable processes (a slight extension of left-continuity of sample paths).

Quadratic Variation, Quadratic Covariation. We shall need to extend quadratic variation and quadratic covariation to stochastic integrals. The quadratic variation of  $I_t(X) = \int_0^t X(u)dW(u)$  is  $\int_0^t X(u)^2 du$ . This is proved in the same way as the case  $X \equiv 1$ , that W has quadratic variation process t. More generally, if  $Z(t) = \int_0^t X(u)dM(u)$  for a continuous martingale integrator M, then  $\langle Z \rangle(t) = \int_0^t X^2(u)d\langle M \rangle(u)$ . Similarly (or by polarization), if  $Z_i(t) = \int_0^t X_i(u)dM_i(u)$  (i = 1, 2),  $\langle Z_1, Z_2 \rangle(t) = \int_0^t X_1(u)X_2(u)d\langle M_1, M_2 \rangle(u)$ .

## Semi-martingales.

It turns out that semi-martingales give the natural class of stochastic integrators: one can define the stochastic integral

$$\int_{0}^{t} H(u)dX(u) = \int_{0}^{t} H(u)dM(u) + \int_{0}^{t} H(u)dA(u)$$

for predictable integrands H (as above), and for semi-martingale integrators X – but for no larger class of integrators, if one is to preserve reasonable convergence and approximation properties for the operation of stochastic integration. For details, see e.g. [P].

With integrands as general as above, stochastic integrals are no longer martingales in general, but only *local martingales* (see e.g. [P]: martingales on each  $[0, T_n]$ , for some sequence of stopping times  $T_n \uparrow \infty$ ). For our purposes, one loses little by thinking of bounded integrands (recall that we usually have a finite time horizon T, the expiry time of an option, and that bounded processes are locally integrable, but not integrable in general).

## 2. Itô's Lemma

Suppose that b is adapted and locally integrable (so  $\int_0^t b(s)ds$  is defined as an ordinary integral, as in Ch. I), and  $\sigma$  is adapted and measurable with  $\int_0^t E(\sigma(u)^2)du < \infty$  for all t (so  $\int_0^t \sigma(s)dW(s)$  is defined as a stochastic integral, as above). Then

$$X(t) := x_0 + \int_0^t b(s)ds + \int_0^t \sigma(s)dW(s)$$

defines a stochastic process X with  $X(0) = x_0$  (which is often called an Itô

process). It is customary, and convenient, to express such an equation symbolically in differential form, in terms of the stochastic differential equation

$$dX(t) = b(t)dt + \sigma(t)dW(t), \qquad X(0) = x_0.$$
 (\*)

Now suppose  $f : \mathbf{R} \to \mathbf{R}$  is of class  $C^2$ . The question arises of giving a meaning to the stochastic differential df(X(t)) of the process f(X(t)), and finding it. Given a partition  $\mathcal{P}$  of [0, t], i.e.  $0 = t_0 < t_1 < \ldots < t_n = t$ , we can use Taylor's formula to obtain

$$f(X(t)) - f(X(0)) = \sum_{k=0}^{n-1} f(X(t_{k+1})) - f(X(t_k))$$
$$= \sum_{k=0}^{n-1} f'(X(t_k)) \Delta X(t_k) + \frac{1}{2} \sum_{k=0}^{n-1} f''(X(t_k) + \theta_k \Delta X(t_k)) (\Delta X(t_k))^2$$

with  $0 < \theta_k < 1$ . We know that  $\sum (\Delta X(t_k))^2 \rightarrow \langle X \rangle(t)$  in probability (so, taking a subsequence, with probability one), and a little more effort gives

$$\sum_{k=0}^{n-1} f''(X(t_k) + \theta_k \Delta X(t_k)) (\Delta X(t_k))^2 \to \int_0^t f''(X(u)) d\langle X \rangle(u).$$

The first sum is easily recognized as an approximating sequence of a stochastic integral (compare the example above), giving

$$\sum_{k=0}^{n-1} f'(X(t_k)) \Delta X(t_k) \to \int_0^t f'(X(u)) dX(u) :$$

**Theorem (Basic Itô Formula)**. If X has stochastic differential given by (\*) and  $f \in C^2$ , then f(X) has stochastic differential

$$df(X(t)) = f'(X(t))dX(t) + \frac{1}{2}f''(X(t))d\langle X\rangle(t),$$

or writing out the integrals,

$$f(X(t)) = f(x_0) + \int_0^t f'(X(u))dX(u) + \frac{1}{2}\int_0^t f''(X(u))d\langle X \rangle(u).$$

More generally, suppose that  $f : \mathbf{R}^2 \to \mathbf{R}$  is a function, continuously differentiable once in its first argument (which will denote time), and twice in

its second argument (space):  $f \in C^{1,2}$ . By the Taylor expansion of a smooth function of several variables we get for t close to  $t_0$  (we use subscripts to denote partial derivatives:  $f_t := \partial f / \partial t$ ,  $f_{tx} := \partial^2 f / \partial t \partial x$ ):

$$f(t, X(t)) = f(t_0, X(t_0)) + (t - t_0)f_t(t_0, X(t_0)) + (X(t) - X(t_0))f_x(t_0, X(t_0)) + \frac{1}{2}(t - t_0)^2 f_{tt}(t_0, X(t_0)) + \frac{1}{2}(X(t) - X(t_0))^2 f_{xx}(t_0, X(t_0)) + (t - t_0)(X(t) - X(t_0))f_{tx}(t_0, X(t_0)) + \dots,$$

which may be written symbolically as

$$df = f_t dt + f_x dX + \frac{1}{2} f_{tt} (dt)^2 + f_{tx} dt dX + \frac{1}{2} f_{xx} (dX)^2 + \dots$$

In this, we substitute  $dX(t) = b(t)dt + \sigma(t)dW(t)$  from above, to obtain

$$df = f_t dt + f_x (bdt + \sigma dW) + \frac{1}{2} f_{tt} (dt)^2 + f_{tx} dt (bdt + \sigma dW) + \frac{1}{2} f_{xx} (bdt + \sigma dW)^2 + \dots$$

Now using the formal multiplication rules  $dt \cdot dt = 0$ ,  $dt \cdot dW = 0$ ,  $dW \cdot dW = dt$  (which are just shorthand for the corresponding properties of the quadratic variations), we expand

$$(bdt + \sigma dW)^2 = \sigma^2 dt + 2b\sigma dt dW + b^2 (dt)^2 = \sigma^2 dt + higher-order terms$$

to get finally

$$df = \left(f_t + bf_x + \frac{1}{2}\sigma^2 f_{xx}\right)dt + \sigma f_x dW + \text{higher-order terms.}$$

As above, the higher-order terms are irrelevant, and summarizing, we obtain  $It\hat{o}$ 's lemma, the analogue for the Itô or stochastic calculus of the chain rule for ordinary (Newton-Leibniz) calculus:

**Theorem (Itô's Lemma)**. If X(t) has stochastic differential given by (\*), then f = f(t, X(t)) has stochastic differential

$$df = \left(f_t + bf_x + \frac{1}{2}\sigma^2 f_{xx}\right)dt + \sigma f_x dW.$$

That is, writing  $f_0$  for  $f(0, x_0)$ , the initial value of f,

$$f = f_0 + \int_0^t (f_t + bf_x + \frac{1}{2}\sigma^2 f_{xx})dt + \int_0^t \sigma f_x dW.$$