

---

# M4A32: Vortex Dynamics

## Problem Sheet 1

---

1. Show that the vorticity equation for the vorticity  $\underline{\omega}$  of a barotropic fluid of density  $\rho$  (so that the fluid pressure  $p = p(\rho)$ ) is given by

$$\frac{D}{Dt} \left( \frac{\underline{\omega}}{\rho} \right) = \frac{\underline{\omega}}{\rho} \cdot \nabla \underline{u}$$

2. By modifying the proof in the case of an ideal fluid, prove that Kelvin's circulation theorem also holds for a barotropic fluid where  $p = p(\rho)$ .

3. Consider the special case of *axisymmetric flow without swirl* in which the velocity field has the special form

$$\underline{u}(r, z, t) = (u_r(r, z, t), 0, u_z(r, z, t))$$

in *cylindrical* polar coordinates. Show that the vorticity vector has the form

$$\underline{\omega}(r, z, t) = (0, \omega(r, z, t), 0)$$

and show that the equation for the scalar function  $\omega(r, z, t)$  is

$$\frac{D}{Dt} \left( \frac{\omega}{r} \right) = 0$$

4. In *spherical* polar coordinates, an incompressible axisymmetric flow without swirl has the form

$$\underline{u}(r, \theta, t) = (u_r(r, \theta, t), u_\theta(r, \theta, t), 0)$$

Define a streamfunction  $\Psi$  for this flow and establish that the vorticity  $\underline{\omega}$  has the form

$$\underline{\omega}(r, \theta, t) = (0, 0, \omega(r, \theta, t))$$

where

$$\omega(r, \theta, t) = -\frac{1}{r \sin \theta} \left( \frac{\partial^2 \Psi}{\partial r^2} + \frac{1}{r^2} \frac{\partial^2 \Psi}{\partial \theta^2} - \frac{\cot \theta}{r^2} \frac{\partial \Psi}{\partial \theta} \right)$$

Hence, by seeking separable solutions, determine the streamfunction  $\Psi$  associated with steady *irrotational* flow of speed  $U$  past a sphere of radius  $a$ .

5. Using the method of Fourier transforms, verify that the three-dimensional solution of

$$\nabla^2 \psi = m \delta(\mathbf{x} - \mathbf{x}_0)$$

which decays at infinity is given by

$$\psi = -\frac{m}{4\pi r}$$

where  $r = |\mathbf{x} - \mathbf{x}_0|$ .

**6.** For an unbounded flow in three dimensions which vanishes at infinity, the Biot-Savart integral is given by

$$\underline{u}(\mathbf{x}) = -\frac{1}{4\pi} \int_{Flow} \frac{1}{r^3} (\mathbf{x} - \mathbf{x}') \wedge \underline{\omega}(\mathbf{x}') dV(\mathbf{x}')$$

where  $r = |\mathbf{x} - \mathbf{x}'|$ . Using this as a starting point, derive the Biot-Savart integral for an unbounded two-dimensional flow (vanishing at infinity) in the plane  $z = \text{constant}$ .

*Hint:* Standard tables show that

$$\int_{-\infty}^{\infty} (x^2 + y^2 + z^2)^{-3/2} dz = \frac{2}{x^2 + y^2}$$